

Oxidative stress and changes in swimming performances at zebrafish model (*Danio Rerio* H. 1822) produced by acute exposure to deltamethrin

Strungaru S.A.¹; Radojković P.²; Dumitru G.³, Nicoara M.³; Plavan G.I.^{3*}; Todirascu-Ciornea E.³

Received: June 218

Accepted: November 2018

Abstract

Delamethrin can easy reach in the aquatic ecosystems and affects the life forms. It produces significant damage to the fish community. It is less know how the fish community reacts in the first hours when insecticide deltamethrin reaches their ecosystem and what biochemical mechanisms of protection are activated. The measurements of the behavior parameters from the tests can explain how fish community reacts before lethal and sublethal deltamethrin dosage. This can be similar to a shock wave produced during an ecological catastrophe where the fish community is involved. Which is the fate of the survivors? Can be their swimming performances recovered to assure their surviving? The aim of this study was to assess on zebrafish model the oxidative stress intensity (MDA, CAT, SOD, GPx) triggered by the acute exposure to lethal and sublethal relevant concentrations of delthametrin and swimming performances recovery of the survivors for: total distance swim, swim velocity, active swim, clockwise rotations, counterclockwise rotations and maximum acceleration.

Keywords: Zebrafish, Deltamethrin, Acute exposure, Oxidative stress, Behavior

1-Faculty of Biology, Department of Research, "Alexandru Ioan Cuza" University of Iasi, Bd. Carol I, 20A, 700505, Iasi, Romania

2-University of Niš, Faculty of Sciences and Mathematics, Department of Biology and Ecology, Višegradska br. 33, 18000 Niš, Republic of Serbia, njorkap@gmail.com

3-Department of Biology, Faculty of Biology, "Alexandru Ioan Cuza" University of Iasi, Bd. Carol I, 20A, 700505, Iasi, Romania

*Corresponding author's Email:gabriel.plavan@uaic.ro

Introduction

The rapid expansion of human population requires more and more water, food and other resources for everyday surviving. Man has been developed new ways to increase the food production by reducing the competition with pests (Chromcova *et al.*, 2015, Fiorino *et al.*, 2018, Gulhan *et al.*, 2012, Gulhan *et al.*, 2014, Orun *et al.*, 2013, Velmurugan *et al.*, 2009). The use of the chemical synthesis compounds is one of the most frequently method applied to control the pests. The synthetic pyrethroids were considered to have a high insecticidal activity, a low toxicity in mammals and no residue in the biosphere (He *et al.*, 1989). At least this was mention in the past. In now days the scientific community agreed that synthetic pyrethroids are quite dangerous in environment to many vertebrate and invertebrate organisms.

Deltamethrin (DM) [(S) α -cyano-3-phenoxybenzyl-(1R)-cis-3-(2,2-dibromovinyl)-2,2dimethylcyclopropane-carboxylate] is a type II pyrethroid extensively used in agriculture and forestry because of high activity against insect pests (Dinu *et al.*, 2010, Dubey *et al.*, 2013, He *et al.*, 1989, Köprücü *et al.*, 2008). This insecticide once released in environment can easy reach in the aquatic ecosystems and it will do damages in life forms populations. It produces significant damage to the fish community. This was proved in various experiments performed under

laboratory conditions with freshwater fish species like: rainbow trout (Aksakal *et al.*, 2010, Velisek *et al.*, 2009), mirror carp (Çalta *et al.*, 2004), *Carassius auratus* L. 1758 (Dinu *et al.*, 2010), common carp (Cengiz, 2006, Köprücü *et al.*, 2004, Svobodová *et al.*, 2003), Nile tilapia (Dubey *et al.*, 2013, Yildirim *et al.*, 2006), zebrafish (DeMicco *et al.*, 2010, Huang *et al.*, 2014), guppy (Israel Stalin *et al.*, 2008, Güneş *et al.*, 2011, Viran *et al.*, 2003), North African catfish (Datta *et al.*, 2003), liver catfish (Kumar *et al.*, 1999), *Pangasius hypophthalmus* S. 1878 (Hedayati *et al.*, 2014), *Ancistrus multispinis* (Da Silva de Assis *et al.*, 2009). It is less know how the fish community reacts in the first hours when insecticide deltamethrin reaches their ecosystem and what biochemical mechanisms of protection will be activated.

Propolis may serve as an antitoxic agent against pesticide toxicity to aquatic animals. It may improve some biochemical markers associated with oxidative stress in fish brain, after exposure to synthetic compounds (Kakoolaki *et al.*, 2013, Orun *et al.*, 2014, Selcuk *et al.*, 2014).

The zebrafish model (*Danio rerio*) is a very popular model organism that is used in scientific research (behavior neuroscience, ecotoxicology, drug development, genotoxicity, brain disorders and ecology). It can be used to measure the pollutants effects upon the fish community and biodiversity (Bartoskova *et al.*, 2013, Fazio *et al.*,

2012, Güneş *et al.*, 2011, Hedayati *et al.*, 2014, Plhalova *et al.*, 2018). The measurements of the behavior parameters from the tests can explain how fish community reacts during the lethal and sublethal acute exposure to deltamethrin which has different intensities. This is similar to a shock wave produced during an ecological catastrophe where the fish community is involved. Which is the fate of the survivors? Can be their swimming performances recovered to assure their surviving?

The aim of this study was to assess on zebrafish model the oxidative stress intensity triggered by the acute exposure to lethal and sublethal relevant concentrations of delthametrin and swimming performances recovery of the survivors.

Materiala and methods

Test Animals

The adults (6-7 months old) of zebrafish with long fin strain were purchased from different breeders to have a higher genetic diversity (Iasi District, Romania). A total number of 120 specimens (60 males and 60 females) were brought in laboratory, housed in two 90 L aquariums and acclimated for one month. During this time, the water was under constant mechanical and biological filtration. It was changed at every 48 h to avoid the intoxication with harmful nitrogen compounds resulted from degradation of organic matter (eg. ammonia and nitrates). The water parameters were

constantly measured with a HI 9828 multiparameter for water analyses from Hanna instruments and in Table 1 are presented the values. The illumination of the aquarium for housing and testing was from white led bands (light intensity was 307.5 LUX) controlled by trigger timers on a 14/10 light/darkness photoperiodic cycle (lights on at 5:00 am and off at 7:00 pm). The fishes were fed with TetraMin Tropical Flakes. Each experimental aquarium had the volume of 7 L, filled up with dechlorinated tap water that was constantly aerated by an air pump. The animals were strictly maintained and treated according to EU Commission Recommendation of 18 June 2007 on guidelines for the accommodation and care of animals used for experimental and other scientific purposes and Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes. This experiment has been approved by the Faculty of Biology from “Alexandru Ioan Cuza” University of Iasi, with the registration number 4465/2017.

Table 1: Water parameters (pH, temperature (t°C), Total Dissolved Solids (TDS), Salinity (Sal.), Conductivity and Oxydation Reduction Potential (ORP)) of the housing aquarium and treatment aquarium.

Housing aquarium					
pH	t°C	TDS (mg L ⁻¹)	Sal.	Conductivity μS cm ⁻¹	ORP
7.66±0.2	24±0.5	276	0.27	553	377
Treatment aquarium					
7.65±0.1	23.5±0.5	274	0.27	552	376

Experimental Design

It was simulated in this study a possible scenario of contamination with deltamethrin in a freshwater ecosystem. In this case we use a homologated insecticide with high quality certificate purchased from market packaged in 2.5 ml vial. The active compound from this insecticide is deltamethrin with the standard certified concentration of 100 g L⁻¹. This was diluted in the treatment aquariums until they were reached the concentrations necessary for this study. Firstly it was studied the scientific literature regarding the concentrations of DM that were used in zebrafish studies. Huang *et al.* (2014) conducted an acute exposure study on adult zebrafish with sublethal concentrations of: 0.15, 1.5, 3.75, 7.5 and 15 µg L⁻¹. The authors studied three behavioral parameters: swimming speed, swimming depth and hyperactivity/surfacing. DeMicco *et al.* (2010) calculated the lethal concentrations (LC) values in zebrafish: LC₅₀ was 40 ± 0.08 µg L⁻¹; 16% mortality at 19 µg L⁻¹ and 84% mortality at 100 µg L⁻¹.

In our study they were selected three relevant concentrations based on the previous studies: 3.12, 12.5 and 25 µg L⁻¹. These were integrated in the scenario presented in this paper. The acute exposure with these concentrations had 72h length. The test animals were separated in four groups with 10 specimens each (5 males and 5 females): group control, group exposed to 3.12 µg L⁻¹, group exposed to 12.5

µg L⁻¹ and group exposed to 25 µg L⁻¹. The acute exposure with these concentrations was repeated for three times to assure the quality control of the statistic tests. The control group simulated a 72 h length exposure to have relevant results about the unharmed fish population.

The behavioral measurements were performed in a multipurpose cross maze that was turned into a T maze. This maze was made from transparent Plexiglas. The infrared light had a source located under it. The images were recorded by an infrared camera connected to a computer and analysed by the software EthoVision XT 11.5 (NOLDUS, Holland) that was previously calibrated for the swimming performance test. The length of each trial was 4 minutes per individual specimen. During this time, each one should explore the area of aquarium. The camera recorded them and the software calculated the raw data for the follow swimming performance variables: total distance swim (cm), swim velocity (cm s⁻¹), active swim (s), clockwise rotations, counterclockwise rotations, maximum acceleration (cm s⁻²). Each group of fish required 24 h for accommodation with experimental aquarium before exposure. In the next day they were conducted pre-treatment tests to establish the unaffected normal behavior of each group. This test was conducted in triplicates for each group with one day before the treatment. In the next day they were administrated the treatments for each experimental

aquarium. In the control aquariums it was not administrated DM. The experiment started at 6 am. The behavioral tests were conducted at different hours (h) during DM exposure and treatment simulation: 2 h, 9 h, 24 h, 48 h and 72 h.

The fishes that presented symptoms similar with paralysis, moments before death were removed from experimental aquariums and killed via head impact. This procedure was applied to all surviving fishes at the end of exposure and treatment simulation. They were stored in labelled plastic bags at - 20°C for oxidative stress analyses.

Oxidative stress analyses

There were measured enzymes involved in the oxidative stress like: superoxide dismutase (SOD), malondialdehyde (MDA), catalase (CAT) and glutathione peroxidase (GPx). Higher or lower activity of these enzymes in the exposed groups compared with the control group represented the biochemical answer deltamethrin toxicity. It was studied in this paper if the selected concentrations can rapidly trigger or not the oxidative stress as biochemical response. The oxidative stress measurements were applied for quantification of pollution on fish community in experimental conditions (Burgos-Aceves *et al.*, 2018, Dinu *et al.*, 2010, Faggio *et al.*, 2016, Fazio *et al.*, 2012, García-Medina *et al.*, 2010, Gobi *et al.*, 2018, Güneş *et al.*, 2011, Lushchak, 2016).

Each frozen fish was weighted and rapidly grinded for oxidative stress analysis. The superoxide dismutase was measured with SOD Assay Kit (Merck, Darmstadt, Germany). The unit of activity for SOD (USOD) was expressed per mg total protein.

The activity of catalase was estimated using the Sinha method (Sinha, 1972), based on colorimetric determination of chromic acetate resulted in the reaction. The wavelength used for determination was set at 570 nm. A unit of measurement (UCAT) represented the enzyme quantity necessary for decomposing of one micromole of hydrogen peroxide in one minute at 20°C and a pH 7. This was expressed per mg total protein.

The glutathione peroxidase activity was spectrophotometric measured at the wavelength of 412 nm. One unit of GPx (UGPx) represented the enzyme quantity necessary to oxidize a μmol GSH per minute. It was measured with Glutathione Assay Kit (Merck, Darmstadt, Germany). This was expressed per mg total protein. The protein concentrations were measured by the Coomassie blue method using bovine serum albumin as a standard (Bradford, 1976).

Malondialdehyde was measured (nM MDA) with Lipid Peroxidation Assay Kit (Merck, Darmstadt, Germany). All the enzymes were spectrophotometric measured with Specord 210 Plus from Analytik Jena, Germany.

Statistical analysis

The one-way ANOVA test followed by the Tukey HSD test has been performed, after the Shapiro-Wilk test, to demonstrate the significant differences variance of the investigated behavioral variables from initial condition of the subjects (pre-treatment) to the end of the treatment in the case of each group. For oxidative stress were compared the differences between control group and treatments groups at the end of acute exposure (ANOVA test followed by the Tukey HSD). All results were presented for this study as average \pm SD. The statistical analyses were carried using OriginPro v.9.3 (2016) software designed and created by OriginLab Corporation, USA.

Results and discussions

According to the published literature the concentration of DM 25 $\mu\text{g L}^{-1}$ and 12.5 $\mu\text{g L}^{-1}$ are below LC_{50} ($40\pm 0.08 \mu\text{g L}^{-1}$) described by DeMicco *et al.*, (2010) and $\text{LC}_{50-24\text{h}}$ ($14.43\pm 1.031 \mu\text{g L}^{-1}$) experimented by Huang *et al.*, (2014). In our experiment, the mortality observed for DM 25 $\mu\text{g L}^{-1}$ and 12.5 $\mu\text{g L}^{-1}$ was 100% in both cases before 9 h exposure time. It was observed the first case of mortality only after 30 minutes exposure time for the concentration 25 $\mu\text{g L}^{-1}$ and 45 minutes exposure time for 12.5 $\mu\text{g L}^{-1}$. Between 10-15 minutes before the death, the zebrafish swim close to the water surface with rapid movements up and down. In the next stage, the fish were semi-paralysed in vertical position in aquarium. Every

time when they touched the bottom of aquarium, they tried to swim with spasmodic moments close to the surface in a corkscrew manner. Moments later it was observed a total paralysis when the fishes lay down to the bottom of aquarium and hyperventilated their gills. From this moment it was just a matter of time till the death. At all death specimens they were observed red spots close to the pectoral and caudal fins. Huang *et al.* (2014) did the follow observations before death of the zebrafish during the acute exposure to DM: rapid gill movement, erratic swimming, swimming in a corkscrew manner, rapid opercular movement and fish remained vertically in water or lay on one side on the bottom of the aquarium moments before death. The authors observed these symptoms for concentrations above 7 $\mu\text{g L}^{-1}$. In our study we observed these symptoms also for 25% of the population exposed to 3.2 $\mu\text{g L}^{-1}$ of DM after 24 h. The fish with these symptoms were excluded from the trial tests of swimming performances. It resulted that DM from the studied insecticide was more toxic than the regular standard solution used in other studies. This can be explained by presence of additives that increased the toxicity level. Anyway the sublethal concentration was 3.2 $\mu\text{g L}^{-1}$.

In Fig. 1 are presented the main picture of the results obtained after the swimming performance tests of the zebrafish without lethal symptoms. In the pre-treatment it was observed homogeneity of the groups regarding

the exploration of the maze area. The modifications started 2 h after acute

exposure with the concentrations $25 \mu\text{g L}^{-1}$ and $12.5 \mu\text{g L}^{-1}$.




Figure 1: The activity of the groups before and after DM exposure with treatment simulation of the control group.

It was observed same pattern between these two groups. They were observed similar patterns between the control group and exposed group with $3.2 \mu\text{g L}^{-1}$.

Swimming performance results

The variable total distance swim shows the capacity of zebrafish to explore the area of the maze. This is similar in a real ecosystem with the capacity of moving away from a pollution source. In the studied groups were observed significant differences (Fig. 2) between

the initial condition of the fish and exposure to high concentrations of DM. The group treated with $25 \mu\text{g L}^{-1}$ had explored in pre-treatment 791.6 ± 264.9 cm per trial and after 2 h exposure time the distance significantly decreased to 337.9 ± 218.6 cm. Similar results were for $12.5 \mu\text{g L}^{-1}$ where in pre-treatment the distance was 721.3 ± 259.7 cm and after 2 h exposure time significantly decreased to 251.5 ± 137 cm. At the fish without symptoms exposed to $3.2 \mu\text{g L}^{-1}$ were not observed significant

differences ($p>0.05$ ANOVA) between pre-treatment (749.6 ± 270.1 cm) and acute exposure at: 2 h (569.64 ± 352

cm), 9h (644.4 ± 405.7 cm), 24 h (505.2 ± 127.9 cm), 48 h (461.5 ± 163.2 cm) and 72 h (426.8 ± 186.1 cm).




Figure 2: Total distance swim and velocity presented as average±SD with the results of one-way ANOVA test.

For the control group were not observed significant differences between pre-treatment (625.9 ± 294.7 cm) and treatment simulation where at 72 h was measured 751.3 ± 190.6 cm.

The variable swim velocity (Fig. 2) was significant decreased at group exposed to $25 \mu\text{g L}^{-1}$ where in pre-treatment the zebrafish had $3.2\pm1.2 \text{ cm s}^{-2}$ and at 2h after exposure was $1.06\pm0.56 \text{ cm s}^{-2}$. Same results were obtained at group exposed to $12.5 \mu\text{g L}^{-1}$ where in pre-treatment had $3.03\pm1.07 \text{ cm s}^{-2}$ and decreased to $1.06\pm0.56 \text{ cm s}^{-2}$. The fish group without symptoms

exposed to $3.12 \mu\text{g L}^{-1}$ had no significant difference ($p>0.05$ ANOVA) between pre-treatment ($3.15\pm1.13 \text{ cm s}^{-2}$) and the end of exposure time at 72h ($1.78\pm0.77 \text{ cm s}^{-2}$). In the control group they were not observed significant differences ($p>0.05$ ANOVA) between pre-treatment ($2.62\pm1.23 \text{ cm s}^{-2}$) and treatment simulation at 72h ($3.14\pm0.8 \text{ cm s}^{-2}$).

Huang *et al.*, (2014) studied same variable during a 360 minutes exposure. The swimming speed in the lowest studied concentration ($0.15 \mu\text{g L}^{-1}$) was increased from the average 39.6 mm s^{-1}

to 49.7 mm s^{-1} . It increased in the highest DM concentration ($15 \mu\text{g L}^{-1}$) from 43.0 mm s^{-1} to 54.0 mm s^{-1} but this happened in the first 60-120 minutes then decreased similar to our results observed at higher concentration. At concentration of $3.75 \mu\text{g L}^{-1}$ the swimming speed significantly decreased at 180-240 min ($22 \pm 5.5 \text{ mm s}^{-1}$) during exposure compared with time interval 0-60 min ($48.7 \pm 3.4 \text{ mm s}^{-1}$). The authors explained that the speed dramatically changed during the DM exposure. In our study at $3.12 \mu\text{g L}^{-1}$ it was not observed any significant changes in the swimming speed. One of the explanations is represented by testing of the zebrafish that had no symptoms.

The variable active swimming (Fig. 3) was relevant to measure response to DM exposure on activity of zebrafish (active/stationary). For the active swim were measure significant differences for the groups exposure to $25 \mu\text{g L}^{-1}$ (between pre-treatment with $215.7 \pm 35.17 \text{ s}$ and 2h with $157.17 \pm 57.79 \text{ s}$) and $12.5 \mu\text{g L}^{-1}$ (between pre-treatment with $216.3 \pm 49.2 \text{ s}$ and 2h with $159.4 \pm 57.7 \text{ s}$).

These results proved that zebrafish become less active similar to a lethargic behaviour at 2h after exposure to these concentrations. In case of the group exposed to $3.12 \mu\text{g L}^{-1}$ were not measured significant differences between activity in pre-treatment ($218.9 \pm 46.9 \text{ s}$) and at the end of exposure at 72h ($162.9 \pm 52.7 \text{ s}$). As for the control group the results of activity

were not significant different between pre-treatment ($191.5 \pm 72.4 \text{ s}$) and the end of simulation at 72h ($224.6 \pm 35.7 \text{ s}$).

The variable maximum acceleration was not affected by DM exposure in all studied cases (Fig. 3). They were not resulted significant differences for all exposed groups. The zebrafish still had the capacity to swim fast on short distance. The values of this variable were for the studied groups: $25 \mu\text{g L}^{-1}$ (between pre-treatment with $54.4 \pm 13.06 \text{ cm s}^2$ and 2h with $50.5 \pm 2.02 \text{ cm s}^2$), $12.5 \mu\text{g L}^{-1}$ (between pre-treatment with $54.9 \pm 9 \text{ cm s}^2$ and 2h with $50.9 \pm 4.6 \text{ cm s}^2$), $3.12 \mu\text{g L}^{-1}$ (between pre-treatment with $53.9 \pm 2.6 \text{ cm s}^2$ and 72h with $50.6 \pm 1.5 \text{ cm s}^2$) and control group (between pre-treatment with $53.3 \pm 3.6 \text{ cm s}^2$ and 72h with $52.3 \pm 1.22 \text{ cm s}^2$).

The number of clockwise and counter clockwise rotations is related with the decisions of fishes during the trial tests of changing their direction in exploration (Fig. 4). Same as in case of the other variables the clockwise rotations significantly decreased at the groups exposed to $25 \mu\text{g L}^{-1}$ DM (between pre-treatment with 5.25 ± 4.2 and 2h with 1.87 ± 1.12) and $12.5 \mu\text{g L}^{-1}$ (between pre-treatment with 4.73 ± 3.53 and 2h with 2 ± 1.6). For the group exposed with $3.2 \mu\text{g L}^{-1}$ were not measured significant differences for clockwise rotations between pre-treatment (5.7 ± 3.3) and 72h (1.3 ± 1.9). Same results were for the control group between pre-treatment (3.7 ± 2.4) and 72h (5.7 ± 3.1). There were significant differences in the groups exposed to 25

$\mu\text{g L}^{-1}$ DM (between pre-treatment with 5.13 ± 2.2 and 2h with 3 ± 3) and $12.5 \mu\text{g L}^{-1}$ (between pre-treatment with 6.7 ± 4.3 and 2h with 1.6 ± 0.9) where this variable significantly decreased. The

group exposed with $3.2 \mu\text{g L}^{-1}$ and the control had not significant variations of this variable.




Figure 3: Active swimming and stationary time presented as average \pm SD with the results of one-way ANOVA test.

Oxidative stress results

The scientific literature was focused on deltamethrin effects produced on the oxidative stress on molluscs, fish and rat model. Köprücü *et al.* (2008) acute exposed (1, 24, 48, 72 and 96 h) the freshwater mussel *Unio elongatulus eucirrus* to DM with different concentrations: 25, 50, 100, 200, 400, 800 and $1,600 \mu\text{g L}^{-1}$. The study concluded that glutathione and catalase levels in digestive gland and gill of freshwater mussel decreased as a response to DM toxicity.

Dinu *et al.* (2010) studied the toxicological effects of a single dose of DM ($2 \mu\text{g L}^{-1}$) on *Carassius auratus gibelio* for 14 days. Their study had been concluded that DM induced oxidative stress in liver and intestine. Dubey *et al.* (2013) proved on their study focused on rat model that in the group treated with deltamethrin, the activity of SOD and CAT significantly decreased.

In our study it was measured the oxidative stress (an imbalance between the production of free radicals and the body's antioxidant defence system) was

triggered or not by the acute exposure to DM at zebrafish in 72 h (Fig. 5). It explained in this scenario if the fish community was capable or not to defend on the studied concentrations. The fish with intoxication symptoms

were collected moments before death and the survivors were sacrificed at the end of the experiment.




Figure 4: Clockwise and counterclockwise rotations presented as average±SD with the results of one-way ANOVA test.

Malondialdehyde activity significantly increased ($*p<0.05$) in the group exposed with $25 \mu\text{g L}^{-1}$ (303.07 ± 16.03 nM MDA) and $12.5 \mu\text{g L}^{-1}$ (281.6 ± 14.88 nM MDA), based on Tukey HSD results, compared with control group (224.7 ± 25.94 nM MDA). There were not significant differences between the group exposed to $3.12 \mu\text{g L}^{-1}$ (232.39 ± 61.9 nM MDA) and control group. MDA is an important reactive metabolite and a marker of LPO. It represents one of the most frequent reactions caused by free radical attack

on biological structures (Dubey *et al.*, 2013).

Superoxide dismutase activity significantly varied between the studied groups ($**p<0.01$ ANOVA). In our studied it was a different pattern in case of the groups exposed to lethal concentrations. The group exposed to $25 \mu\text{g L}^{-1}$ had the lowest values (0.22 ± 0.04 USOD/mg protein) and was significant lower than control (0.35 ± 0.06 USOD/mg protein), group treated with $3.12 \mu\text{g L}^{-1}$ (0.43 ± 0.01 USOD/mg protein) and group treated with $12.5 \mu\text{g L}^{-1}$ (0.7 ± 0.04 USOD/mg

protein). This suggested a significant oxidative stress caused on the zebrafish body in a short period of time. Its body is fighting against the DM toxicity. On the other hand, the group treated with $12.5 \mu\text{g L}^{-1}$ had values of SOD

significant higher than control group suggesting that at this concentration the antioxidant biochemical equipment of the fish has problems.




Figure 5: The analysis of the average \pm SD with the results of one-way ANOVA test.

Between the group treated with the lowest concentration and control there were not significant differences but the trend suggests that these DM concentrations had antioxidant effect similar with the group $12.5 \mu\text{g L}^{-1}$. Stress conditions result in the decrease in antioxidant enzyme activity because to their excessive utilization (Dubey *et al.*, 2013).

The catalase activity showed an excessive utilization of the antioxidant enzymes promoting the oxidative stress that increased with the concentration of DM dissolved in water. Fish

biochemical system is able to fight from the first moments against the DM acute toxicity. The variations of CAT activity were significant lower between control group (97.9 ± 13.2 UCAT/mg protein) < 3.12 $\mu\text{g L}^{-1}$ (72.9 ± 10.3 UCAT/mg protein); 12.5 $\mu\text{g L}^{-1}$ (64.3 ± 3.2 UCAT/mg protein) < 25 $\mu\text{g L}^{-1}$ (34.3 ± 7.1 UCAT/mg protein).

Glutathione peroxidase was significant lower at the group exposed to 25 $\mu\text{g L}^{-1}$ (0.04 ± 0.008 UGPx/mg protein) compared with the others. Between the other groups did not result significant differences concerning the

activity of this enzyme: control (0.06 ± 0.007 UGPx/mg protein), $3.12 \mu\text{g L}^{-1}$ (0.07 ± 0.003 UGPx/mg protein) and $12.5 \mu\text{g L}^{-1}$ (0.06 ± 0.008 UGPx/mg protein).

Based on these results DM trigger the oxidative stress based on the activity of these enzymes. It is possible that at the zebrafish that survived at the lowest concentration of DM, the oxidative stress was decreasing as a slow recovery against its toxicity.

Conclusions

The concentrations of deltamethrin used in this study were more toxic than similar used in other studies. This can be explained by other chemicals additives added in the insecticide that increased its toxicity. The scenario presented in our study showed that the fishes which survived behind the shock wave of DM toxicity are still capable to same swimming performances as the control group. This may help them to swim away from the pollution source and to repopulate the habitat after this danger passed. Further experiments are necessary to measure the damages on fish population that survived on acute exposure to DM regarding complex cognitive processes like memory, aggressive behaviour and anxiety. The toxicity of this compound was related with the oxidative stress. In case of lethal concentrations it was triggered fast to keep for a longer time the body functions. For sublethal concentrations, it was also activated the oxidative stress

but with a lower intensity suggesting a possible recovery of the organism.

Acknowledgement

This study was financial supported by the project Resources Pilot Center for Cross-Border Preservation of the Aquatic Biodiversity of Prut River MIS-ETC 1150 Romania – Ukraine – Republic of Moldova. The project PN-III-P1-1.2-PCCDI-2017-0245 SUSTENVPRO was also acknowledged.

References

- Aksakal, E., Ceyhun, S.B., Erdoğlan, O. and Ekinci, D., 2010. Acute and long-term genotoxicity of deltamethrin to insulin-like growth factors and growth hormone in rainbow trout. *Comp Biochem Physiol.*, Part C, 152(4), 451-455.
- Bartoskova, M., Dobsikova, R., Stancova, V., Zivna, D., Blahova, J., Marsalek, P., Zelnickova, L., Bartos, M., Di Tocco, F.C. and Faggio, C., 2013. Evaluation of ibuprofen toxicity for zebrafish (*Danio rerio*) targeting on selected biomarkers of oxidative stress. *Neuro Endocrinology Letters*, 34, 102-108.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), 248-254.
- Burgos-Aceves, M.A., Cohen, A., Smith, Y. and Faggio, C., 2018.

- MicroRNAs and their role on fish oxidative stress during xenobiotic environmental exposures. *Ecotoxicology and Environmental Safety*, 148, 995-1000.
- Çalta, M. and Ural, M.Ş., 2004.** Acute toxicity of the synthetic pyrethroid deltamethrin to young mirror carp, *Cyprinus carpio*. *Fresenius Environmental Bulletin*, 13(11a), 1179-1183.
- Cengiz, E.I., 2006.** Gill and kidney histopathology in the freshwater fish *Cyprinus carpio* after acute exposure to deltamethrin. *Environmental Toxicology and Pharmacology*, 22(2), 200–204.
- Chromcova, L., Blahova, J., Zivna, D., Plhalova, L., Casuscelli Di Tocco, F., Divisova, L., Prokes, M., Faggio, C., Tichy, F. and Svobodova, Z., 2015.** NeemAzal T/S – toxicity to early-life stages of common carp (*Cyprinus carpio* L.). *Veterinarni Medicina*, 60(1), 23–30.
- Commission Recommendation, 2007.** Guidelines for the accommodation and care of animals used for experimental and other scientific purposes (notified under document number C (2007) 2525). *Official Journal of the European Union*, Vol. 50.
- Da Silva de Assis, H.C., Nicareta, L., Salvo, L.M., Klemz, C., Truppel, J.H. and Calegari, R., 2009.** Biochemical Biomarkers of Exposure to Deltamethrin in Freshwater Fish, *Ancistrus multispinis*. *Brazilian Archives of Biology and Technology*, 52 (6), 1401-1407.
- Datta, M. and Kaviraj, A., 2003.** Acute toxicity of the synthetic pyrethroid deltamethrin to freshwater catfish *Clarias gariepinus*. *Bulletin of Environmental Contamination and Toxicology*, 70(2), 296–299.
- DeMicco, A., Cooper, K.R., Richardson, J.R. and Lori White, A., 2010.** Developmental neurotoxicity of pyrethroid insecticides in zebrafish embryos. *Toxicological Sciences*, 113(1), 177–186.
- Dinu, D., Marinescu, D., Munteanu, M.C., Staicu, A.C., Costache, M. and Dinischiotu, A., 2010.** Modulatory effects of deltamethrin on antioxidant defense mechanisms and lipid peroxidation in *Carassius auratus gibelio* liver and intestine. *Archives of Environmental Contamination and Toxicology*, 58(3), 757–764.
- Directive of the European Parliament and of the Council, 2010.** Protection of animals used for scientific purposes. *Official Journal of the European Union*, Vol. 53.
- Dubey, N., Khan, A.M. and Raina, R., 2013.** Sub-acute Deltamethrin and Fluoride Toxicity Induced Hepatic Oxidative Stress and Biochemical Alterations in Rats. *Bulletin of Environmental Contamination and Toxicology*, 91(3), 334–338.

- Faggio, C., Pagano, M., Alampi, R., Vazzana, I. and Felice, M.R., 2016.** Cytotoxicity, haemolymphatic parameters and oxidative stress following exposure to sub-lethal concentrations of quaternium-15 in *Mytilus galloprovincialis*. *Aquatic Toxicology*, 180, 258-265.
- Fazio, F., Faggio, C., Marafioti, S., Torre, A., Sanfilippo, M. and Piccione, G., 2012.** Comparative study of haematological profile on *Gobius niger* in two different habitat sites: Faro Lake and Tyrrhenian Sea. *Cahiers de Biologie Marine*, 53, 213-219.
- Fiorino, E., Sehonova, P., Plhalova, L., Blahova, J., Svobodova, Z. and Faggio, C., 2018.** Effect of glyphosate on early life stages: comparison between *Cyprinus carpio* and *Danio rerio*. *Environmental Science and Pollution Research (international)*, 25(9), 8542–8549.
- García-Medina, S., Razo-Estrada, A.C., Gómez-Oliván, L.M., Amaya-Chávez, A., Madrigal-Bujaidar, E. and Galar-Martínez, M., 2010.** Aluminum-induced oxidative stress in lymphocytes of common carp (*Cyprinus carpio*). *Fish Physiology and Biochemistry*, 36(4), 875–882.
- Gobi, N., Vaseeharan, B., Rekha, R., Vijayakumar, S. and Faggio, C., 2018.** Bioaccumulation, cytotoxicity and oxidative stress of the acute exposure selenium in *Oreochromis mossambicus*. *Ecotoxicology and Environmental Safety*, 162, 147-159.
- Gulhan, M.F., Duran, A., Selamoglu Talas, Z., Kakoolaki, S. and Mansouri, S.M., 2012.** Effects of Propolis on microbiologic and biochemical parameters of Rainbow trout (*Oncorhynchus mykiss*) after exposure to the pesticide. *Iranian Journal of Fisheries Sciences*, 11(3), 490-503.
- Gulhan, M.F., Selamoglu Talas, Z., Erdogan, K. and Orun, I., 2014.** The Effect of Propolis on Gill, Liver, Muscle Tissues of Rainbow Trout (*Oncorhynchus mykiss*) Exposed to Various Concentrations of Cypermethrin. *Iranian Journal of Fisheries Sciences*, 13(3), 684-701.
- Güneş, E. and Yerli, S.V., 2011.** Effects of deltamethrin on lipase activity in guppies (*Poecilia reticulata*). *Turkish Journal of Fisheries and Aquatic Sciences*, 11, 473-476.
- He, F., Wang, S., Liu, L., Chen, S., Zhang, Z. and Sun, J., 1989.** Clinical manifestations and diagnosis of acute pyrethroid poisoning. *Archives of Toxicology*, 63(1), 54-58.
- Hedayati, A. and Tarkhani, R., 2014.** Hematological and gill histopathological changes in iridescent shark, *Pangasius hypophthalmus* (Sauvage, 1878) exposed to sublethal diazinon and deltamethrin concentrations. *Fish Physiology and Biochemistry*, 40(3), 715–720.

- Huang, Y., Zhang, J., Han, X. and Huang, T., 2014.** The use of zebrafish (*Danio rerio*) behavioral responses in identifying sublethal exposures to deltamethrin. *International Journal of Environmental Research and Public Health*, 11(4), 3650-3660.
- Israel Stalin, S., Kiruba, S. and Sam Manohar Das, S., 2008.** A comparative study on the toxicity of a synthetic pyrethroid, deltamethrin and a neem based pesticide, azadirachtin to *Poecilia reticulata*, Peters 1859 (Cyprinodontiformes: Poeciliidae). *Turkish Journal of Fisheries and Aquatic Sciences*, 8(1), 01-05.
- Kakoolaki, S., Selamoglu Talas, Z., Cakir, O., Ciftci, O. and Ozdemir, I., 2013.** Role of Propolis on Oxidative Stress in Fish Brain. *Basic Clin Neurosc*, 4(2), 45-50.
- Köprücü, K. and Aydin, R., 2004.** The toxic effects of pyrethroid deltamethrin on the common carp (*Cyprinus carpio* L.) embryos and larvae. *Pesticide Biochemistry and Physiology*, 80(1), 47-53.
- Köprücü, K., Yonar, E. and Seker, E., 2008.** Effects of deltamethrin on antioxidant status and oxidative stress biomarkers in freshwater mussel, *Unio elongatulus eucirrus*. *Bulletin of Environmental Contamination and Toxicology*, 81(3), 253-257.
- Kumar, S., Lata, S. and Gopal, K., 1999.** Deltamethrin induced physiological changes in freshwater cat fish *Heteropneustes fossilis*. *Bulletin of Environmental Contamination and Toxicology*, 62(3), 254-258.
- Lushchak, V.I., 2016.** Contaminant-induced oxidative stress in fish: a mechanistic approach. *Fish Physiology and Biochemistry*, 42(2), 711-747.
- Orun, I., Dogru, M.I., Erdogan, K., Dogru, A., Ongun, A., Yuksel, E. and Talas-Selamoğlu, Z., 2013.** Effects of acute and chronic exposure to glyphosate on common carp (*Cyprinus carpio* L.) hematological parameters: The beneficial effect of propolis. *Fresenius Environmental Bulletin*, 22(9) 2504-2509.
- Orun, I., Selamoglu Talas, Z., Gulhan, M.F. and Erdogan, K., 2014.** Role of propolis on biochemical and hematological parameters of *Oncorhynchus mykiss* exposed to cypermethrin. *Survey in Fisheries Sciences.*, 1(1), 21-35.
- Plhalova, L., Blahova, J., Divisova, L., Enevova, V., Casuscelli Di Tocco, F., Faggio, C., Tichy, F., Vecerek, V. and Svobodova, Z., 2018.** The effects of subchronic exposure to NeemAzal T/S on zebrafish (*Danio rerio*). *Chemistry and Ecology.*, 34 (3), 199-210.
- Selcuk Aldemir, O., Selamoglu Talas, Z., Gulhan Fuat, M., Cakir, O., Dastan Durna, S. and Dogan, H., 2014.** Role of Propolis on Oxidative Stress in Various Tissues of Fish.

- Fresenius Environmental Bulletin*, 23 (12c), 3546 – 3550.
- Sinha, A.K., 1972.** Colorimetric assay of catalase. *Analytical Biochemistry*, 47(2), 389-394.
- Strungaru, S.A., Robea, M.A., Plavan, G., Todirascu-Ciornea, E., Ciobica, A. and Nicoara, M., 2018.** Acute exposure to methylmercury chloride induces fast changes in swimming performance, cognitive processes and oxidative stress of zebrafish (*Danio rerio*) as reference model for fish community. *Journal of Trace Elements in Medicine and Biology*, 47, 115-123.
- Svobodová, Z., Lusková, V., Drastichová, J., Svoboda, M. and Îlábek, V., 2003.** Effect of deltamethrin on haematological indices of common carp (*Cyprinus carpio* L.). *Acta Veterinaria Brno*, 72, 79–85.
- Velisek, J., Svobodova Z. and Piackova, V., 2009.** Effects of acute exposure to bifenthrin on some haematological, biochemical and histopathological parameters of rainbow trout (*Oncorhynchus mykiss*). *Veterinarni Medicina - Czech*, 54(3), 131–137.
- Velmurugan, B., Selvanayagam, M., Cengiz, E.I. and Unlu, E., 2009.** Histopathological changes in the gill and liver tissues of freshwater fish, *Cirrhinus mrigala* exposed to dichlorvos. *Brazilian Archives of Biology and Technology*, 52 (5), 1291-1296.
- Viran, R., Erkoç, F.Ü., Polat, H. and Koçak, O., 2003.** Investigation of acute toxicity of deltamethrin on guppies (*Poecilia reticulata*). *Ecotoxicology and Environmental Safety*, 55(1), 82–85.
- Yildirim, M.Z., Benli, A.Ç.K., Selvi, M., Özkul, A., Figen, E. and Koçak, O., 2006.** Acute toxicity, behavioral changes, and histopathological effects of deltamethrin on tissues (gills, liver, brain, spleen, kidney, muscle, skin) of Nile tilapia (*Oreochromis niloticus* L.) fingerlings. *Environmental Toxicology*, 21(6), 541–622.