

## Toxic metal contamination and total organic carbon content in the meat of the main fish species imported and sold in Romanian's supermarkets

Strungaru S.A.<sup>1,4</sup>; Nicoara M.<sup>2,4\*</sup>; Gorban C.F.<sup>3</sup>; Paduraru E.<sup>4</sup>; Plavan G.I.<sup>2</sup>

Received: June 2019

Accepted: September 2019

### Abstract

In Romania as in Western Europe in the last years a mass panic installed among the consumers of cheap fish meat imported mostly from Vietnam – *Pangasianodon hypophthalmus* (S 1878). This problem was intensively fuelled by mass media and different movies uploaded on internet. The metal compounds, once released in the environment, can easily contaminate the food webs and finally the body of the highest rank consumers. The toxic metals in high concentrations are damaging the biological functions and in the end they may trigger the death of the organism (depending on the concentration and the body weight). These metals cannot be biodegraded in the environment; they are quickly transferred from the environment and accumulated in the biological tissues. This study investigated the level of arsenic, mercury, lead and cadmium in the meat of *Pangasianodon hypophthalmus*, *Merluccius hubbsi*, *Scomber scombrus*, from nine different brands using HR-CS GF-AAS and HG-AAS techniques.

**Keywords:** Toxic metals, Total organic carbon, HR-CS GF-AAS, HG-AAS

---

1-Alexandru Ioan Cuza” University of Iasi, Department of Research, Faculty of Biology, 700505 Iasi, Romania

2-Alexandru Ioan Cuza” University of Iasi, Faculty of Biology, Department of Biology, 700505 Iasi, Romania

3-University of Agricultural Sciences and Veterinary Medicine “Ion Ionescu de la Brad”, Iasi, Aleea Mihail Sadoveanu nr. 3, 700490, Romania

4-Alexandru Ioan Cuza” University of Iasi, Doctoral School of Geosciences, Faculty of Geography-Geology, Bd. Carol I, 700505, Iasi, Romania

\*Corresponding author's Email: mirmag@uaic.ro

## Introduction

Food resource contamination with various toxic compounds is an actual problem for the entire human society (Chaid *et al.*, 2014; Yildirim *et al.*, 2009; Manthey-Karl *et al.*, 2016; Guimarães *et al.*, 2016; Vilavert *et al.*, 2017; Perugini *et al.*, 2014; Madrid *et al.*, 2016; Plavan *et al.*, 2017; Strybgary *et al.*, 2015), being associated with severe health problems and death. Fish meat is one of the cheapest animal proteins that can be found on market in high amounts. There are two types of fish resources from where the consumer can chose to buy: fish produced by aquaculture industry and wild captured fish. Both of these resources are under high anthropogenic pressure. For instance, the aquaculture industry is involving a series of chemicals harmful for human health (aquatic pollutants, antibiotics, hormones and artificial fish food). The environmental pollution and the industrial fishing that damage the aquatic ecosystems are playing the major role in case of the wild fish resource. The meat of panga fish became popular among the consumer thanks to its low price, texture, no smell and merchandising form into filets with no skin and spine (Ruiz-de-Cenzano *et al.*, 2013, Ionescu *et al.*, 2016). In most of the cases, the regular consumer is influenced by the product price and not by quality. Panga fish meat requires high industrial processing techniques (Orban *et al.*, 2008) compared with wild captured fish like *Merluccius hubbsi* (Marini, 1933) and *Scomber scombrus* (Linnaeus, 1758). The toxic

metals resulted from anthropogenic activities can damage the terrestrial and aquatic biota because they are accumulating very fast, they are toxic and they are not biodegradable (Ionescu *et al.*, 2016; Caunii *et al.*, 2015; Micu *et al.*, 2016). The development of analytical methods, and technology significantly improved the quality and speed of detection for toxic contaminants in food. Mercury, arsenic, cadmium and lead are the most toxic metals that can threat the human population if occur in high amounts.

In this study we investigated the traces of mercury, arsenic, cadmium and lead in meat of *Pangasianodon hypophthalmus* (panga fish), *Scomber scombrus* (Atlantic mackerel) and *Merluccius hubbsi* (Argentine hake) from nine brands with different origins, common in the Romanian supermarkets, using the methods HR-CS GF-AAS and HG-AAS. Total organic carbon content was measured for each brand, this being related to the fish meat quality.

## Experimental part

The fish was purchased from three different supermarkets located in the City of Iasi, Romania. These are the largest from this city and belong to same franchise that has same supermarket brand all around Romania. Because of the globalization same product that belongs to a brand can be found everywhere. Each supermarket may have different providers for the same product. In this study it was purchased from each supermarket once

per month from February to April a bag of frozen fish or 1 kg self-weighting bag for each studied species: *Pangasianodon hypophthalmus*, *Merluccius hubbsi* and *Scomber scombrus*. These products are the cheapest fish from the market. In the supermarkets we found for same species different brands. The panga fish was found only in industrial processed frozen fillets. Atlantic mackerel and Argentine hake were originated from captured oceanic fish and they were commercialized frozen with or without guts.

#### *Sample preparation*

The samples were transported in laboratory and prepared according to the following protocol: fish purchased every month was unfrosted and the water excess was removed using clean paper towels; from each brand it was sampled with PE tools a mass of 10 g of muscle per sample (in triplicates  $n=3$ ) that was well homogenized in an agate mortar with pestle; from this homogenate was weighted a mass of 1 g in triplicates that were inserted in TFM vessels and mixed for digestion with nitric acid (4 mL, 65% Suprapur Merck) and hydrogen peroxide (1 mL, Merck) (Plavan *et al.*, 2017; Strungaru *et al.*, 2015) after 25 minutes of reaction the vessels were inserted in Speedwave MWS-2 produced by Berghof using a specific program (Plavan *et al.*, 2017; Strungaru *et al.*, 2015); the resulted liquid samples were transferred into 50 mL volumetric flasks (Brand) and diluted with ultrapure water produced

from double distilled water; for calibration were used certified standard solutions from Merck for arsenic (As), mercury (Hg), lead (Pb) and cadmium (Cd) with each concentration of 1000 mg L<sup>-1</sup>; mercury was measured with cold vapour hydride generator coupled to AAS that required a sodium borohydride solution and hydrochloric acid (3M, Merck). For the total organic carbon measurements the meat homogenate was dried at 75°C and grinded again to a fine powder. This was analysed in triplicates for each sample, using the direct combustion at 1300°C in solid module coupled to Focus Radiation NDIR detector, multi N/C 2100S produced by AnalytikJena Germany.

#### *Validation of method and results*

A total number of 243 samples were prepared and analysed in this study for the metals content. We used fish muscle dry powder certified by European Reference Material ERM-BB422 from Joint Research Centre Institute for Reference Materials and Measurements to validate the method optimization and the results. We prepared 6 samples from this material by weighting a mass of 0.3 g that followed same protocol as a normal sample. Arsenic, cadmium and lead were measured with HR-CS GF-AAS model ContrAA 600 produced by AnalytikJena Germany, which was optimized for this sample matrix. Mercury was measured with cold vapour hydride generator coupled to AAS model Avanta produced by GBC Australia. Argon 5.0 (99.99%) was used

as carrier gas. The results for validation obtained after the analysis of the reference material were: As – certified value of  $12.7 \pm 0.7 \mu\text{g g}^{-1}$  and obtained value of  $11.5 \pm 0.5 \mu\text{g g}^{-1}$ ; Hg - certified value of  $0.601 \pm 0.03 \mu\text{g g}^{-1}$  and obtained value of  $0.65 \pm 0.4 \mu\text{g g}^{-1}$ ; Cd – certified value of  $0.0075 \pm 0.0018 \mu\text{g g}^{-1}$  and obtained value of  $0.0081 \pm 0.0012 \mu\text{g g}^{-1}$ ; Pb – was not presented in standard certificate and for our method was below LOD.

#### *Statistical analysis*

Shapiro-Wilk normality test was performed for the data sets. One-way ANOVA followed by the Tukey HSD test was conducted to demonstrate the significant variance of toxic metal between the brands and fish species (Plavan *et al.*, 2017; Strungaru, 2015). All the statistical analyses were carried with OriginPro v.9.3 (2016) software created by OriginLab Corporation, USA.


#### **Results and discussion**

In Figure 1 are presented the values of arsenic reported as average  $\pm$  95% confidence for each brand and species. There were significant differences between the brands for same species. There were not observed significant differences between the species regarding the concentration of this element ( $p > 0.05$ ). Panga fish had significant variation between brands ( $*p < 0.05$ ), with the highest values for Brand\_2 but with no significant differences compared with other species. The confidence interval is very

large suggesting that not all the fishes are originated from same aquaculture system. The concentrations were: Brand\_1 with  $1.38 \pm 1.79 \mu\text{g g}^{-1}$ , Brand\_2 with  $2.33 \pm 2.98 \mu\text{g g}^{-1}$  and Brand\_3 with  $1.55 \pm 2.01 \mu\text{g g}^{-1}$ .

The Argentine hake had significant variations between the brands ( $*p < 0.05$ ) with large confidence intervals of values suggesting that they may be mixed with fish from different catchment areas exposed to pollution sources, possible close to the shore. The concentrations were: Brand\_4 with  $2.17 \pm 2.95 \mu\text{g g}^{-1}$ , Brand\_5 with  $2.2 \pm 2.86 \mu\text{g g}^{-1}$  and Brand\_6 with  $1.32 \pm 1.87 \mu\text{g g}^{-1}$ . For the Atlantic mackerel no differences between brands were observed ( $p > 0.05$ ). The concentrations were: Brand\_7 with  $2.01 \pm 2.55 \mu\text{g g}^{-1}$ , Brand\_8 with  $1.91 \pm 2.47 \mu\text{g g}^{-1}$  and Brand\_9 with  $1.73 \pm 2.58 \mu\text{g g}^{-1}$ . The general concentrations expressed as average  $\pm$  SD for each studied species for arsenic were: panga fish  $1.79 \pm 0.61 \mu\text{g g}^{-1}$ , Argentine hake  $1.91 \pm 0.67 \mu\text{g g}^{-1}$  and Atlantic mackerel  $1.88 \pm 0.53 \mu\text{g g}^{-1}$ . This element is not included in (EC) No 1881/2006 EU legislation that is setting maximum levels for certain contaminants in foodstuffs. In this case was not proved if the concentrations of this element exceeded the maximum permitted values.

Mercury had no significant variance between the brands grouped for each species.




**Figure 1: Concentrations of As from each studied brand and species, reported as average $\pm$ 95% confidence with the measurement unit expressed as  $\mu\text{g g}^{-1}$  wet weight. The species with significant variance for the one-way ANOVA are marked with line and  $*-p<0.05$ .**

There were significant differences ( $*p<0.05$ ), instead, between panga fish and the wild species captured (Fig. 2).

Panga fish had no significant differences ( $p>0.05$ ) between brands, with the values: Brand\_1 with  $0.27\pm 0.34 \mu\text{g g}^{-1}$ , Brand\_2 with  $0.22\pm 0.28 \mu\text{g g}^{-1}$  and Brand\_3 with  $0.25\pm 0.33 \mu\text{g g}^{-1}$ . Similar to arsenic, the confidence interval is large. Argentine hake had no significant differences ( $p>0.05$ ) between brands, with the values: Brand\_4 with  $0.14\pm 0.19 \mu\text{g g}^{-1}$ , Brand\_5 with  $0.19\pm 0.22 \mu\text{g g}^{-1}$  and Brand\_6 with  $0.17\pm 0.30 \mu\text{g g}^{-1}$ . Same differences were for Atlantic mackerel: Brand\_7 with  $0.32\pm 0.32 \mu\text{g g}^{-1}$ , Brand\_8 with  $0.17\pm 0.23 \mu\text{g g}^{-1}$  and Brand\_9 with  $0.17\pm 0.23 \mu\text{g g}^{-1}$ . The general comparisons between the species provide significant results ( $*p<0.05$ ). The highest concentration of total mercury was measured in panga

fish:  $0.25\pm 0.06 \mu\text{g g}^{-1}$  (average $\pm$ SD) that was significant higher than in Argentine hake with  $0.17\pm 0.06 \mu\text{g g}^{-1}$  and Atlantic mackerel with  $0.19\pm 0.06 \mu\text{g g}^{-1}$ . This element is included in (EC) No 1881/2006 with the maximum value of  $0.50 \mu\text{g g}^{-1}$  ( $\text{mg kg}^{-1}$ ) allowed for fishery products and fish muscle. The panga fish did not exceed the limit level for mercury in average but for some samples the levels are at the maximum accepted limit. There were doubts about the quality of the meat according to this element. No significance difference between the two wild species originated from oceanic capture was observed.

The concentrations for cadmium and lead with very low traces were measured in the domain of nanograms. For cadmium and lead were not recorded significant differences between the brands grouped for same species ( $p>0.05$ ).

Hg  $\mu\text{g g}^{-1}$ 


**Figure 2: Concentrations of Hg from each studied brand and species, reported as average $\pm$ 95% confidence with the measurement unit expressed as  $\mu\text{g g}^{-1}$  wet weight. The comparisons between species with significant variance for Tukey test are marked with  $*-p<0.05$ .**

Differences were observed, instead, between panga fish, Argentine hake and Atlantic mackerel in the case of cadmium (Fig. 3).

The values for cadmium reported as average $\pm$ 95% confidence were: Brand\_1 with  $0.34\pm 0.62$   $\text{ng g}^{-1}$ , Brand\_2 with  $0.27\pm 0.44$   $\text{ng g}^{-1}$ , Brand\_3 with  $0.21\pm 0.28$   $\text{ng g}^{-1}$ , Brand\_4 with  $2.9\pm 3.46$   $\text{ng g}^{-1}$ , Brand\_5 with  $1.78\pm 2.11$   $\text{ng g}^{-1}$ , Brand\_6 with  $2.97\pm 4.43$   $\text{ng g}^{-1}$ , Brand\_7 with  $5.64\pm 6.86$   $\text{ng g}^{-1}$ , Brand\_8 with  $3.77\pm 4.9$   $\text{ng g}^{-1}$ , Brand\_9 with  $4.1\pm 6.82$   $\text{ng g}^{-1}$ . The highest concentration of total cadmium was measured in Atlantic mackerel with  $4.46\pm 1.61$   $\text{ng g}^{-1}$  (average $\pm$ SD), significantly higher than in Argentine hake with  $2.49\pm 0.89$   $\text{ng g}^{-1}$  and panga fish with  $0.27\pm 0.16$   $\text{ng g}^{-1}$ . This element is included in (EC) No

1881/2006 with the maximum value of  $100$   $\text{ng g}^{-1}$  ( $0.1$   $\text{mg kg}^{-1}$ ) allowed for fishery products and fish muscle. This element was in the ultra-trace domain proving that the products were not contaminated. In case of panga fish the exposure was significantly lower than in the two wild species captured.

Lead had no significant differences ( $p>0.05$ ) between brands and measured samples. The values for lead reported as average $\pm$ 95% confidence were: Brand\_1 with  $3.65\pm 6.03$   $\text{ng g}^{-1}$ , Brand\_2 with  $5.7\pm 6.7$   $\text{ng g}^{-1}$ , Brand\_3 with  $3.9\pm 5.4$   $\text{ng g}^{-1}$ , Brand\_4 with  $5.9\pm 10.3$   $\text{ng g}^{-1}$ , Brand\_5 with  $3.1\pm 5.8$   $\text{ng g}^{-1}$ , Brand\_6 with  $4.1\pm 5.7$   $\text{ng g}^{-1}$ , Brand\_7 with  $3.59\pm 7.2$   $\text{ng g}^{-1}$ , Brand\_8 with  $6.35\pm 9.04$   $\text{ng g}^{-1}$ , Brand\_9 with  $5.85\pm 7.48$   $\text{ng g}^{-1}$ .




**Figure 3: Concentrations of Cd from each studied brand and species, reported as average $\pm$ 95% confidence with the measurement unit expressed as  $\mu\text{g g}^{-1}$  wet weight. The comparisons between species with significant variance for Tukey test are marked with  $*-p<0.05$ .**

The highest concentration of total lead was measured in Atlantic mackerel with  $5.33\pm 2.53 \text{ ng g}^{-1}$  (average $\pm$ SD) that was not significantly higher ( $p>0.05$ ) than in Argentine hake with  $4.32\pm 2.72 \text{ ng g}^{-1}$  and panga fish with  $4.52\pm 1.55 \text{ ng g}^{-1}$ . This element is included in (EC) No 1881/2006 with the maximum value of  $300 \text{ ng g}^{-1}$  ( $0.3 \text{ mg kg}^{-1}$ ) allowed for fishery products and fish muscle. This element was in the ultra-trace domain proving that the products were not contaminated (Fig 4).

The total organic carbon (TOC) content was reported as average $\pm$ SD for dry muscle. Between the brands of panga fish there were significant differences: Brand\_1 with  $393.7\pm 38.2 \text{ g kg}^{-1}$ , Brand\_2 with  $518 \pm 23.5 \text{ g kg}^{-1}$  and Brand\_3 with  $328 \pm 37.1 \text{ g kg}^{-1}$ . The high values in Brand\_2 raise doubts

about the quality. High values of TOC may be caused by a high content of fatty acids and other organic carbon compounds. The Argentine hake had significant differences between the brands: Brand\_4 with  $197.8\pm 18.6 \text{ g kg}^{-1}$ , Brand\_5 with  $475.3 \pm 11.2 \text{ g kg}^{-1}$  and Brand\_6 with  $403.6 \pm 40.8 \text{ g kg}^{-1}$ . The explanation for the low content in the Brand\_4 may consist of the poor food resources and the fishing period when the fish was losing in weight. There were not observed significant differences for TOC in Atlantic mackerel: Brand\_7 with  $438.1\pm 16.1 \text{ g kg}^{-1}$ , Brand\_8 with  $405.9 \pm 21.8 \text{ g kg}^{-1}$  and Brand\_9 with  $468.8 \pm 11.2 \text{ g kg}^{-1}$ . It is known that this fish is one of the richest in fatty acids and omega 3.

**Pb** ng g<sup>-1</sup>

**Figure 4: Concentrations of Cd from each studied brand and species, reported as average±95% confidence with the measurement unit expressed as  $\mu\text{g g}^{-1}$  wet weight.**

The similar values obtained at this species proved that the quality was the same for all purchased brands.

### Conclusions

The toxic metals did not exceed the maximum permitted values, according to EU legislation, in fish meat of the studied brands excepting the panga fish meat where the mercury concentrations in some samples reached the maximum legal limit. This raised doubts regarding the quality of the meat provided by certain brands. Cadmium and lead were in the domain of ultra-trace elements with harmless low concentrations. The Atlantic mackerel was the only species from this study with no differences between the brands regarding the toxic metal content and TOC. The consumer should have more information about the levels of possible contaminants from

food to choose himself the level of exposure. From the price point of view the consumer can pay more for a quality product with less contaminants because it may require more investments. It will be not right to pay more for a low quality only to bring more profit to the sellers. This should be controlled by government authorities in any country.

### Acknowledgments

This study was financial supported by the project Resources pilot centre for cross-border preservation of the aquatic biodiversity of Prut River MIS-ETC 1150 Romania – Ukraine – Republic of Moldova

### Conflict of interest

Authors declare that they have no competing interests.

## References

- Caunii, A., Negrea, A., Pentea, M., Samfira, I., Motoc, M., Butnariu, M., 2015.** Mobility of Heavy Metals from Soil in the Two Species of the Aromatic Plants, *Revista de Chimie* (Bucharest), 66, 3, pp. 382-386.
- Chaid, A., Hilali, M., Benlhachimi, A., Bouzid, T., 2014.** Mobility of Heavy Metals from Soil in the Two Species of the Aromatic Plants. *Food Chemistry*, 147, pp. 357-360.
- Commision Regulation (EC), 2006.** No 1881/2006 setting maximum levels for certain contaminants in foodstuffs. Section 3: Metals 364/18-21; Section 6: Polycyclic aromatic hydrocarbons 364/22-24. *Official Journal of the European Union*.
- Guimarães, C.F.M., Mársico, E.T., Monteiro, M.L.G., Lemos, M., Mano, S.B., Conte Jr, C. A. C., 2016.** The chemical quality of frozen Vietnamese *Pangasius hypophthalmus* fillets. *Food Science and Nutrition*, 4, 3, pp. 398-408.
- Ionescu, P., Deak, G., Diacu, E., Radu, V.M., 2016.** Assessment of heavy metal levels in water, sediment and fish from Plumbuita Lake, Romania. *Revista de Chimie* (Bucharest), 67, 11, 2016, pp. 2148-2150.
- Madrid, E., Gil, F., García, M., Debenedetti, A.L., Trelis, M., Fuentes, M.V., 2016.** Potential risk analysis of human anisakiasis through the consumption of mackerel, *Scomber scombrus*, sold at Spanish supermarket *Food Control*, 66, pp. 300-305.
- Manthey-Karl, M., Lehmann, I., Ostermeyer, U., Schröder, U., 2016.** Meat Composition and Quality Assessment of King Scallops (*Pecten maximus*) and Frozen Atlantic Sea Scallops (*Placopecten magellanicus*) on a Retail Level. *Food*, 5, 58, pp. 1-14.
- Micu, L.M., Petanec, D.I., Iosub-Ciur, M.D., Andrian, S., Popovici, R.A., Porumb, A., 2016.** The heavy metals content in leaves of the forest fruits (*Hippophae rhamnoides* and *Rubus fruticosus*) from the tailings dumps mining. *Revista de Chimie* (Bucharest), 67, 1, pp. 64-68.
- Orban, E., Nevigato, T., Di Lena, G., Masci, M., Casini, I., Gambelli, L., Caproni R., 2008.** New trends in the seafood market. Sutchi catfish (*Pangasius hypophthalmus*) fillets from Vietnam: Nutritional quality and safety aspects. *Food Chemistry*, 110, pp. 383-389.
- Perugini, M., Visiano, P., Manera, M., Zaccaroni, A., Olivieri, V., Amorena, M., 2014.** Heavy metals (As, Cd, Hg, Pb, Cu, Zn, Se) concentration in muscle and bone of four commercial fish caught in the central Adriatic Sea, Italy. *Environmental Monitoring Assessment*, 186, pp. 2205-2213.
- Plavan, G., Jitar, O., Teodosiu, C., Nicoara, M., Micu, D., Strungaru, S.A., 2017.** Toxic metals in tissues of fishes from the Black Sea and associated human health risk exposure. *Environmental Science and Pollution Research*, 24, no.8, pp. 7776-7787

- Ruiz-de-Cenzano, M., Beser, U., Cervera, M.L., De La Guardia, M.F., 2013.** Fast determination of fish mineral profile. Application to Vietnamese pangasid fish. *Ecotoxicology and Environmental Safety*, 95, pp. 195-201.
- Strungaru, S.A., Nicoara, M., Jitar, O., Plavan, G., 2015.** Influence of urban activity in modifying water parameters, concentration and uptake of heavy metals in *Typha latifolia* L. into a river that crosses an industrial city. *Journal of Environmental Health Science & Engineering*, 13, 5, pp.1-11.
- Strungaru, S.A., Nicoara, M., Rau, M.A., Plavan, G., Micu, D., 2015.** Do you like to eat fish? An overview of the benefits of fish consumption and risk of mercury poisoning. *Analele Ştiinţelor Univ. "Alexandru Ioan Cuza" din Iaşi*, s. Biologie Animală, Tom LXI, pp. 117-123.
- Usydus, Z., Szlinder-Richert, J., Adamczyk, M., Szatkowska, U., 2011.** Marine and farmed fish in the Polish market: Comparison of the nutritional value. *Food Chemistry*, 126, pp. 78-84.
- Vilavert, L., Borrell, F., Nadal, M., Jacobs, S., Minnens, F., Verbeke, W., Marques, A., Domingo, J. L., 2017.** Health risk/benefit information for consumers of fish and shellfish: FishChoice, a new online tool. *Food and Chemical Toxicology*, 104, pp. 79-84.
- Yildirim, Y., Gonulalan, Z., Narin, I., Soylak, M., 2009.** Evaluation of trace heavy metal levels of some fish species sold at retail in Kayseri, Turkey. *Environmental Monitoring and Assessment*, 149, pp. 223-228.