

## Zooplankton of Hancağız Dam Lake (Gaziantep - Turkey)

Saler, S<sup>1\*</sup>; Alıř, N<sup>2</sup>

Received: June 2013- Accepted: January 2014

### Abstract

Zooplankton samples were taken seasonally from Hancağız Dam Lake between April 2012- May 2013 period. During the study, a total of 52 species (34 belong to genus Rotifera, 11 to Cladocera and 7 Copepoda) were found in Hancağız Dam Lake. Based on the number of individuals, rotifers were the dominant group in the dam lake (65.4%) followed by Cladocera (21.2%) and Copepoda (13.4%). All of the zooplanktonic species have been detected for the first time in Hancağız Dam Lake. The datas determined from the zooplankton community in Hancağız Dam Lake were analysed by Shannon Wiener species richness index ( $H^1$ ), Margalef diversity index (D) and Sorenson similarity index.  $H^1$  value was found highest with 3.73 in spring. D index value was calculated in its highest level in spring with 6.31.

**Keywords:** Rotifera, Cladocera, Copepoda, Hancağız Dam Lake, Turkey

---

1,2- Firat University Faculty of Fisheries

\*Corresponding author's email: serapsaler@gmail.com

## Introduction

Zooplanktons occupy the second trophic level in the food chain, the first being occupied by phytoplankton. In lake ecosystems, these organisms are the main food source for invertebrates, fishes, and sometimes for aquatic birds. Some species have been reported as characteristic indicators of water quality and trophic level of lakes (Sladeczek, 1983; Herzig, 1987; Saksena, 1987). Studies on the zooplankton fauna of Turkey have been conducted by Gündüz (1984, 1987, 1997), Ustaoglu (1986, 1989), Dumont and De Ridder (1987), Ustaoglu and Balık (1990), Segers et al. (1992), Akbulut (Emir) (2000), Altındağ and Yiğit (2002), Yiğit (2002), Bekleyen (2003), Ustaoglu (2004), Güher and Kırız (2004), Yıldız et al. (2007), Saler and Haykır (2011), Bulut and Saler (2013 a,b). A detailed study on the zooplankton fauna of Hancağız Dam Lake had not been done before.

A detailed study on the zooplankton fauna of Hancağız Dam Lake had not been done before. This study was done to obtain insight into the composition of the zooplankton

fauna of Hancağız Dam Lake and to contribute to the knowledge of the biological diversity of inland waters in Turkey.

## Material and Methods

Hancağız Dam Lake was located in Nizip district of Gaziantep, was built between 1985 - 1989 for irrigation. Lake volume is 100 hm<sup>3</sup> and area is 4.33 km<sup>2</sup> (URL, 1). For determining the zooplankton of Hancağız Dam Lake, samples were taken from three stations seasonally between April 2012 and May 2013 (Figure 1).

Zooplankton samples were collected with a standard plankton net (Hydrobios 55 µm mesh size) through vertical and horizontal hauls from 3 stations. Samples were preserved in 4% formaldehyde in 250 mL plastic bottles. The species were identified using Scourfield and Harding (1941), Dussart (1969), Flössner (1972), Kolisko (1974), Harding and Smith (1974), Koste (1978a,b), Kiefer (1978), Korovchinsky (1992), Reedy (1994), Einsle (1996).



Figure 1: Hancağız Dam Lake and the location of the stations

pH values were measured with Lamotte (pH 5-WC) pH meter, water temperature and dissolved oxygen values were measured by Oxi 315i/SET oxygen meter in situ.

The number of species in a community is referred to as species richness. We used Shannon species richness index formula.

$$H = -\sum_{i=1}^S (P_i \cdot \ln P_i)$$

where,

H = the Shannon diversity index  
 $P_i$  = fraction of the entire population made up of species  $i$  (proportion of a species  $i$  relative to total number of species present, not encountered)

S = numbers of species encountered

One of the measures of species diversity is the Margalef Index. It is calculated from the total number of species presented and the abundance or total number of individuals. The higher index means the greater diversity.

$D = (S - 1) / \log_{10} N$  where

D = Margalef Index

S = The number of species

N = Total number of individuals

The widely used Sørensen similarity index measures similarity in species composition for two sites or stations, A and B, by the

$CS = \frac{2ab}{a+b}$ , where

a = is the number of species found in site A

b = is the number of species in site B

ab = is the number of species shared by the two sites (Magurran, 2004).

## Results

In Hancağız Dam Lake a total of 52 taxa, 34 from rotifers, 11 from cladocerans, and 7 from copepods were identified.

Based on the number of individuals, rotifers were the dominant group in the dam lake (65.4%) followed by Cladocera (21.2%) and Copepoda (13.4%). Dominant Rotifera species was observed as *Polyarthra dolichoptera* followed by *Keratella cochlearis*. *Bosmina longirostris*, *Chydorus sphaericus* and *Ceriodaphnia reticulata* were the dominant Cladoceran species. *Cyclops vicinus* and *Thermo cyclops crassus* were the dominant Copepoda species. The seasonal distributions of species are given in table 1.

In Hancağız Dam Lake *A. saltans*, *A. priodonta*, *B. angularis*, *B. bidentata*, *B. quadridentatus*, *C. gibba*, *F. opoliensis*, *H. fennica*, *K. longispina*, *K. cochlearis*, *K. quadrata*, *K. tecta*, *L. bulla*, *L. luna*, *M. trigona*, *P. dolichoptera*, *S. pectinata*, *S. stylata*, *T. patella* from Rotifera; *A. rectangulata*, *B. longirostris*, *C. reticulata*, *C. sphaericus*, *D. galeata* and *D. rostrata* from Cladocera; *A. denticornis*, *C. vicinus*, and *T. crassus* from Copepoda were observed in all seasons. Spring has got the most species richness with 41 species in Station III and 38 species in Station I. The least species richness were recorded in Station III with 7 species and Station I with 11 species in winter. *P. dolichoptera* the dominant rotifers species observed in all seasons in all stations except Station III in winter. *B. longirostris* from Cladocera and *C. vicinus* from Copepoda had got the highest frequency of occurrence in their groups.

**Table 1: Seasonal distribution of zooplankton fauna of Hancığ Dam Lake**

Seasons	Autumn			Winter			Spring			Summer		
Stations	1	2	3	1	2	3	1	2	3	1	2	3
<b>SPECIES</b>												
<b>Rotifera</b>												
<i>Ascomorphasaltans</i> Bartsch, 1870	-	+	+	-	-	+	+	-	+	-	+	+
<i>Asplanchnapriodonta</i> (Gosse, 1850)	-	-	+	-	-	+	+	-	-	+	+	-
<i>Brachionusangularis</i> Gosse, 1851	-	+	+	-	-	+	+	+	+	-	+	-
<i>Brachionusbidentata</i> (Anderson, 1889)	+	+	-	-	+	-	+	+	+	-	-	+
<i>Brachionuscaudatus</i> (Barrois& Daday,1894)	+	+	-	-	-	-	+	+	-	-	-	-
<i>Brachionusquadridentatus</i> Hermann, 1783	+	-	+	+	-	-	+	+	-	+	-	+
<i>Cephalodellagibba</i> (Ehrenberg, 1830)	-	-	+	+	-	+	+	-	+	+	-	+
<i>Colurellaadriatica</i> Ehrenberg, 1831	-	+	-	-	+	-	-	+	+	-	-	-
<i>Colurellacolurus</i> (Ehrenberg, 1830)	+	+	-	-	-	-	+	-	+	-	+	-
<i>Euclanisdilatata</i> Ehrenberg, 1832	-	-	-	-	+	-	+	+	+	-	-	-
<i>Filinalongiseta</i> (Ehrenberg, 1834)	+	+	-	-	+	-	-	-	+	+	-	-
<i>Hexarthrafennica</i> (Levander, 1892)	-	-	-	-	-	-	+	-	+	-	+	+
<i>Hexarthraintermedia</i> Wierzejski, 1929	-	-	-	+	-	-	-	-	-	+	-	+
<i>Kellicottialongispina</i> (Kellicott, 1879)	+	-	+	+	-	-	+	+	+	-	+	-
<i>Keratellacochlearis</i> (Gosse,1851)	+	+	+	-	+	-	+	+	+	+	+	-
<i>Keratellaquadrata</i> (Muller, 1786)	-	+	-	-	-	-	-	+	+	-	-	-
<i>Keratellatecta</i> (Gosse,1851)	+	-	-	+	-	-	+	-	+	-	+	+
<i>Keratellatropica</i> (Apstein, 1907)	-	-	+	-	-	-	+	+	+	-	-	+
<i>Keratellavalga</i> Ehrenberg, 1834	+	-	-	-	-	-	-	+	+	+	+	-
<i>Lecane bulla</i> (Gosse, 1886)	+	-	+	-	+	-	+	-	+	-	-	+
<i>Lecaneclosterocerca</i> (Schmarda, 1859)	-	-	-	-	+	-	+	-	+	-	-	+
<i>Lecaneluna</i> (Muller, 1776)	+	-	+	+	-	-	-	+	-	-	+	-
<i>Mytilinatrigena</i> Harring, 1913	-	+	+	-	+	-	+	+	+	+	-	+
<i>Notholcasquamula</i> (Muller, 1786)	-	+	-	+	+	-	-	-	-	-	-	-
<i>Philodinaroseola</i> Ehrenberg, 1832	-	-	-	-	-	-	-	+	+	+	-	-
<i>Polyarthradolichoptera</i> Idelson, 1925	+	+	+	+	+	-	+	+	+	+	+	+
<i>Pompholyxsulcata</i> Hudson, 1885	-	+	-	-	-	-	-	+	+	-	-	-
<i>Rotarianeptunia</i> (Ehrenberg, 1832)	-	-	-	-	-	-	+	+	+	+	-	+
<i>Rotariarotatoria</i> (Pallas, 1766)	-	-	-	-	-	-	+	-	+	+	-	-
<i>Synchaetapectinata</i> Ehrenberg, 1832	+	-	+	-	+	-	-	+	+	+	-	-
<i>Synchaetastylata</i> Wierzejski, 1893	-	-	+	-	-	-	+	-	+	-	-	+
<i>Testudinella patina</i> Hermann, 1783	+	-	-	-	+	-	+	-	+	-	+	-

<i>Trichocercapucina</i> Wierzejski & Zacharias, 1893	-	+	-	-	-	-	+	+	-	+	-	+
<i>Trichocercasimilis</i> Wierzejski, 1893)	-	-	+	-	-	-	+	+	-	+	+	+
<b>Cladocera</b>												
<i>Alonarectangulata</i> Sars, 1862	+	-	+	+	-	-	+	+	-	+	-	+
<i>Bosminalongirostris</i> (Muller, 1785)	+	+	-	-	+	-	+	+	+	+	+	+
<i>Ceriodaphniapulchella</i> Sars, 1862	+	-	-	-	-	-	-	+	+	-	-	-
<i>Ceriodaphniareticulata</i> (Jurine, 1820)	+	+	-	+	-	+	+	+	-	+	-	+
<i>Chydorusphaericus</i> (O. F. Müller, 1776)	-	+	+	-	-	+	+	+	+	-	+	-
<i>Diaphanosomabirgei</i> Korinec, 1981	-	-	+	-	-	-	+	-	+	-	+	-
<i>Daphniacucullata</i> Sars, 1862	-	-	+	-	-	-	+	+	+	-	-	+
<i>Daphniagaleata</i> Sars, 1865	+	-	-	+	-	-	-	+	+	+	-	-
<i>Daphnialongispina</i> O. F. Müller, 1785	+	+	-	-	-	-	+	-	+	-	-	+
<i>Disparalonarostrata</i> (Koch, 1841)	-	+	-	-	+	-	-	+	-	-	+	-
<i>Leydigialeydigi</i> (Schoedler, 1863)	-	-	+	-	-	-	-	+	+	-	-	-
<b>Copepoda</b>												
<i>Acanthopdiaptomusdenticornis</i> (Wierjesky, 1887)	+	-	+	+	-	-	+	+	+	-	+	-
<i>Cyclopsstrenuus</i> Fisher, 1851	+	+	-	-	+	-	+	+	+	-	+	+
<i>Cyclops vicinus</i> Uljanin, 1875	+	+	+	+	-	+	+	+	+	+	-	+
<i>Diacyclopsbicuspidatus</i> (Claus, 1857)	-	+	+	-	-	-	+	-	+	+	-	+
<i>Megacyclopsviridis</i> (Jurine, 1820)	+	-	-	-	-	-	+	-	+	-	+	-
<i>Nitocrahibernica</i> (Brady, 1880)	-	-	-	-	-	-	+	+	+	-	-	-
<i>Thermocyclopscrassus</i> (Fischer, 1853)	+	+	-	-	+	-	+	+	+	+	+	+

Table 2: Seasonal Sorenson Similarity Index in Hancağız Dam Lake

Seasons and similarity percentages	
Autumn-Winter= 80.00%	Winter-spring= 74.67%
Autumn- spring= 91.48%	Winter –summer=75.67%
Autumn-summer= 85.05%	Spring-summer=92.47%

Table 3: Seasonal variation of Margalef Diversity Index (D) and Shannon Wiener Species Richness Index (H') in Hancağız Dam Lake

Index	Seasons			
	Autumn	Winter	Spring	Summer
D	5.61	4.42	6.31	5.44
H'	3.28	2.89	3.73	3.09

There was a decrease in total zooplankton species richness in winter and a sharp increase in spring and autumn months. The most species were recorded in spring (32 rotifers, 11 copepods, 7 cladoceran, totally 50 species), but the less taxa were observed in winter (23 rotifers, 6 copepods, 4 cladoceran, totally 33 species). According to the stations, the most number of species were

recorded in the third station with 41 species in spring and which the less were also in the third station with 9 species in winter (Table 1). H' value was found highest with 3.73 in spring. D index value was calculated in spring months as 6.3 (Table 3). Water temperature, dissolved oxygen and pH values of the Hancağız Dam Lake were recorded in the field and shown in the table 4.

**Table 4: Seasonal average values of temperature, dissolved oxygen and pH recorded in Hancağız Dam Lake**

Seasons	Autumn			Winter			Spring			Summer		
	1	2	3	1	2	3	1	2	3	1	2	3
Temperature (C°)	15.1	14.8	14.5	9.5	9.1	8.9	16.8	16.3	16.0	24.8	23.9	23.8
Dis. Oxygen(mg/L)	9.1	9.5	10.2	11.3	10.9	9.9	8.6	8.4	8.3	7.4	7.8	7.6
pH	7.2	6.8	6.6	6.9	7.2	7.7	7.6	7.9	8.4	8.9	8.6	8.1

Water temperature values were changed between 8.9-24.8C°, dissolved oxygen 7.4-11.3-mg/L, pH 6.6-8.9.

## Discussion

Zooplankton species are important indicators for aquatic habitats since most of them are used to determine the quality, the trophic level and level of population in lakes and streams. For example, *Keratella cochlearis* and *Polyarthra dolichoptera* species of Rotifera are indicators of productive habitats, while *Notholca squamulais* known as indicators of cold waters (Kolisko, 1974). *K. cochlearis* and *P. dolichoptera* were determined in all seasons and *N. squamula* was observed in cold seasons, in autumn and winter in Hancağız Dam Lake. Kolisko (1974), reported that *P. dolichoptera* and *K. cochlearis* are perennial species while, *Notholca squamulais* a winter form. The results of this study have got a great accordance with her report. Species richness of Rotifera was found quite high when compared to Cladocera and Copepoda in Turkish inland waters (Ustaoglu

and Balık 1990, Bozkurt, 2004, 2006, Dirican and Musul 2008, Gaygusuz et al. 2004, Yiğit and Altındağ 2005, Yıldız et al., 2007, Saler and Haykır 2011, Bulut and Saler 2013 a,b). Parallely to this result, in Hancağız Dam Lake 34 Rotifer a species were found among 52 identified zooplankton species. According to Stember and Gannon (1978) Rotifera forms an important part of biomass in eutrophic water systems. In the present study, Rotifera appeared as dominant group (65.4%).

All the recorded Rotifer species in the present study are widely distributed around the world (Segers, 2007). Also many of the recorded species are common in Turkey (Saksena 1987, Dumont and De Ridder 1987, Kaya and Altındağ, 2007). Only 11 species of Cladocera were observed in Dam Lake. Among the identified species *Leydigia leydigi*, *Disparalonarostrata*, *Diaphanosomabirgei* and

*Ceriodaphniapulchella* were rarely found in Hancağız Dam Lake. *Chydorus sphaericus*, *Bosmina longirostris*, *Ceriodaphnia reticulata* and *Alonare ctangulata* were observed throughout all seasons. *Cyclops vicinus* and *Acanthodiptomus denticornis* were the observed in all seasons but *Nitocra hibernica* was only recorded in spring. *Cyclops vicinus* and *Acanthodiptomus denticornis* are the common Copepod species in Turkey inland water (Ustaoglu, 2004).

According to the results of this study, Rotifera is recorded as the dominant zooplankton group. The number of zooplankton species showed an increase in spring and autumn and decrease in summer and winter. The zooplanktonic fauna structure of Asi River, Kesikköprü Dam Lake, Gelingülü Dam Lake, Keban Dam Lake, Kepektaş Dam Lake, Sürgü Dam Lake, Kalecik Dam Lake, Beyhan Dam Lake and Uzunçayır Dam Lake were showed similarities with our findings (Bozkurt et al. 2002, Yiğit 2006, Kaya and Altındağ 2007, Tellioglu and Akman, 2007, Saler 2009, İpek and Saler 2013 and Bulut and Saler 2013b, 2014, Saler, et al. 2014). In all of these dam lakes, rotifers were found to be the dominant species as species richness and frequency of occurrence. All of these species are recorded for the first time in Hancağız Dam Lake. Bekleyen (2001), recorded thirty four Rotifer species from Devegeçidi Dam Lake. She reported 12 Rotifer species from Brachionidae. In Hancağız Dam Lake 10 species were recorded from Brachionidae. It was reported that species from Brachionidae were represented in high

number of species in eutrophic waters (Kolisko 1974). The ecological features of the recorded species show that most of them are cosmopolitan and littoral inhabiting (Kolisko, 1974). Additionally, among the recorded species, *Bosmina longirostris* and *C. vicinus*, *P. dolichoptera*, *K. cochlearis* are well known indicators of eutrophy (Ryding and Rast, 1989). *P. dolichoptera*, *K. cochlearis* are predominant in the Dam Lake. *Pompholyx sulcata* and *Hexarthra intermedia* were rarely found in the Dam Lake. *Philodina roseola*, *Rotaria neptunia* and *Rotaria rotatoria* were recorded as the organisms of polluted waters (Kolisko, 1974, Edmondson, 1959). These three species were recorded in spring and summer period.

Some of the zooplankton species recorded in Hancağız Dam Lake were known species of eutrophic waters as *A. priodonta*, *B. calyciflorus*, *B. angularis*, *K. quadrata*, *K. cochlearis*, *E. dilatata*, *L. luna*, *F. longiseta*, *P. dolichoptera*, *B. longirostris*, *C. sphaericus*, *A. denticornis*, *C. vicinus*, and *T. crassus*. Some of the Rotifer species as *L. bulla* (oligotrophic), *B. urceolaris*, *B. calyciflorus*, *B. angularis*, *K. quadrata*, *K. cochlearis*, *P. quadricornis*, *E. dilatata*, *L. luna*, *F. longiseta*, *T. cylindrica*, and *T. patina* are also the species of eutrophic waters. *B. longirostris*, *C. sphaericus*, from Cladocera are the organisms of oligotrophic-eutrophic waters (Gündüz, 1984; Saksena, 1987; Ustaoglu, 1989; Kolisko, 1974; Gutiérrez-Aquirre and Suárez-Morales, 2000).

We identified 52 zooplankton species in the lake. Among these species 19 species (11 rotifers, 6 cladocerans, and 2 copepods) informed that the lake's trophic level was in a

mesotrophic state and vulnerable to eutrophication. In conclusion, in terms of zooplankton diversity and species richness indexes values ( $H'$  and  $D$  index values) of Hancağız Dam Lake it could be assumed that the dam lake has got mesotrophic-eutrophic character and thus, candidate to eutrophication.

## References

- Akbulut(Emir), N., 2000.** Community structure of zooplanktonic organisms in Lake Akşehir. *Turkish Journal of Zoology*, 24, 271-278.
- Altındağ, A. and Yiğit, S., 2002.** The zooplankton fauna of Lake Burdur. *Ege Üniversitesi Su Ürünleri Dergisi*, 19 (1-2), 129-132.
- Bekleyen, A. 2001.** A Taxonomical Study on the Rotifera Fauna of Devegeçidi Dam Lake (Diyarbakır-Turkey). *Turkish Journal of Zoology*, 25, 251-255
- Bekleyen, A. 2003.** A Taxonomical study on the Zooplankton of Göksu Dam Lake, (Diyarbakır), *Turkish Journal of Zoology*, 27, 95-100.
- Bozkurt, A., 2004.** Akdeniz Bölgesindeki Bazı Akarsuların Zooplankton (rotifer, kladocerve kopepod) Faunası Üzerine İlk Gözlemler, *Türk Sucul Yaşam Dergisi*, 3, 65-70.
- Bozkurt, A., 2006.** Yenişehir Gölü (Reyhanlı, Hatay) Zooplanktonu. *Ege Üniversitesi Su Ürünleri Dergisi*, 23(1), 39-43.
- Bozkurt, A., Göksu, M. Z. L., Sarıhan, E. And Taşdemir, M., 2002.** Asi Nehri Rotifer Faunası (Hatay, Türkiye). *Ege Üniversitesi Su Ürünleri Dergisi*, 19(1-2), 63-67
- Bulut, H. and Saler, S., 2013a.** Ladik Gölü (Samsun) Zooplanktonu Üzerine İlk Gözlemler, *Su Ürünleri Mühendisleri Dergisi*, pp. 51-54, pp. 74-78
- Bulut, H. and Saler, S., 2013b.** Kalecik Baraj Gölü (Elazığ- Türkiye) Zooplanktonu. *Fırat Üniversitesi Fen Bilimleri Dergisi*, 25(2), 99-103
- Bulut, H. and Saler, S., 2014.** Zooplankton of Beyhan Dam Lake (Elazığ-Turkey). *Turkish Journal of Science & Technology*, 9(1), 23-28
- Dirican, S. and Musul, H., 2008.** Çamlığöze Baraj Gölü (Sivas-Türkiye) Zooplanktonu Faunası Üzerine Bir Çalışma, *Süleyman Demirel Üniversitesi Fen Bilimleri Dergisi*, 12(1), 17-21.
- Dumond, H. J. and De Ridder, M., 1987.** Rotifers from Turkey, *Hydrobiologia*, 147, 65-73.
- Dussart, B., 1969.** Les Copepodes des Eaux Continentales d'Europe Occidentale Tome II Cyclopoïdes et Biologie, N. Boubee et Cie, Paris.
- Edmondson, W. T., 1959.** Fresh Water Biology Second edition, University of Washington, Seattle, 1248P.
- Einsle, U., 1996.** Guides to the Identification of the Microinvertebrates of the Continental Waters of the World, Copepoda: Cyclopoida, Genera Cyclops, Megacyclops, Acanthocyclops, SPB Academic Pub., Germany.
- Flössner, D., 1972.** Krebstiere, Crustacea. Kiemen and Blattfüßer Brachiopoda Fischlaube, Branchiura, Tierwelt-Deutschlands, 60. Tielweb. Gustav Fischer Verlag, Jena, 501P.
- Gaygusuz, Ö., Acıpinar, H., Dorak, Z. and Temel, M., 2004.** İznik Gölü zooplankton



- gruplarının tespitive mevsimselde gelişimi *XVII. Ulusal Biyoloji Kongresi*, pp. 21-24
- Güher, H. and Kırgız, T., 2004.** The Copepoda (Crustacea) freshwater fauna of Turkish Thrace region (Edirne, Kırklareli, Tekirdağ). *Pakistan Journal of Biological Science*, 7(5), 834-837.
- Gündüz, E., 1984.** Karamukve Hoyran Gölleri'nde zooplankton türlerinin tespitive kirlenmenin zooplankton üzerindeki etkisi, *PhD, thesis*, Hacettepe Üniversitesi. Fen Bilimleri Enstitüsü, Ankara, 83P.
- Gündüz, E., 1987.** Karamıkve Hoyran göllerinin cladocera Crustacea) türleri üzerine taksonomik bir çalışma. *Turkish Journal of Zoology*, 11, 26-36.
- Gündüz, E., 1997.** Türkiye İçsularında Yaşayan Cladocera (Crustacea) Türlerinin Listesi. *Turkish Journal of Zoology*, 21, 37-45.
- Gutiérrez-Aguirre, M. and Suárez-Morales, E., 2000.** The Eurasian *Thermocyclops crassus* (Fischer, 1853) (Copepoda, Cyclopoida) found in Southeastern Mexico. *Crustaceana*. 73(6), 705-713.
- Harding, J. P. and Smith W. A., 1974.** A Key to the British Freshwater, Biological Association Scientific Publication, No: 18, Westmorland.
- Herzig, A., 1987.** The analysis of planktonic rotifers populations. A plea for long-term investigations. *Hydrobiologia*, 147, 163-187.
- Ipek, Alış, N., and Saler, S. 2013.** Zooplankton of Sürgü Dam Lake (Malatya- Turkey). *Bitlis Eren Üniversitesi Journal of Science*, 2(1), 40-44.
- Kaya M. and Altındağ, A., 2007.** Zooplankton Fauna and Seasonal changes of Gelingülü Dam Lake (Yozgat, Turkey) *Turk Journal of Zoology*, 31, 347-351.
- Kiefer, F., 1978.** Das Zooplankton der Binnengewässer 2. Teil, Freilebende Copepoda, Die Binnengewässer Band XXVI, Schweizer band'sche Verlag buch hand lung, Stuttgart.
- Kolisko, W. R., 1974.** Planktonic Rotifers Biology and Taxonomy Biological Station. Lunz of The Austrian Academy of Science, Stuttgart.
- Korovchinsky, N. M., 1992.** Sididae and Holopedidae (Crustacea: Daphniiformes), Guides to Identification of the Micro invertebrates of the Continental Waters of the World, SPB Academic Pub., Netherlands.
- Koste, W., 1978 a.** Die Radertiere Mitteleuropas I. Textband, Berlin.
- Koste, W., 1978b.** Die Radertiere Mitteleuropas II. Tafelband, Berlin.
- Maguran, A., 2004.** Measuring biological diversity. Blackwell Publishing; Oxford
- Reedy, R. Y., 1994.** Copepoda: Calanoida: Diaptomidae. SPB Academic Publication, Leiden.
- Ryding, S. O. and Rast. W., 1989.** The Control of Eutrophication of Lakes and Reservoirs, Man Biosphere Series, Parthenon Publication Group, Vol 1, USA.
- Saksena, N. D., 1987.** Rotifers as indicator of water quality, *Hydrobiology*, 15, 481-485.
- Saler, S. and Haykır, H., 2011.** Zooplankton Composition of Pulumur Stream (Tunceli-Turkey), *Journal of Animal and Veterinary Advances*, 10(11), 1401-1403.
- Saler, S., Haykır, H. and Baysal, N., 2014.** Zooplankton of Uzunçayır Dam Lake

- (TunceliI-TURKEY), *Journal of fisheriessciences.com*, 8(1), 1-7
- Segers, H., Emir, N. and Mertens, J., 1992.** Rotifera from North and North east Anatolia (Turkey). *Hydrobiologia*, 245, 179-189.
- Segers, H. 2007.** Annotated checklist of the rotifers (Phylum Rotifera), with notes on nomenclature, taxonomy and distribution. *Zootaxa*, 1564, 104P.
- Scourfield, D. J. and Hardig, J. P., 1966.** Freshwater Biology, Science Publication, New York.
- Sladeczek, V., 1983.** Rotifers as indicators of water quality, *Hydrobiologia*, 100, 169-201.
- Stemberger, R. S. and Gannon, J. E., 1978.** Zooplankton (especially crustaceans and rotifers) as indicators of water quality, *Transactions of the American Microscopical Society*, 97, 16-35.
- Tellioglu, A. and Akman, F., 2007.** A Taxonomical Study on the Rotifera Fauna in Pertek Region of Keban Dam Lake, *Ege Üniversitesi. Journal of Fisheries and Aquatic Sciences*, 24(1-2) 135-136.
- URL1.2014.** <http://www2.dsi.gov.tr/bolge/dsi20/gaziantep.htm>
- Ustaoglu, M. R., 1986.** Zooplankton (metazoa) of the Karagöl (Yamanlar-İzmir-Türkey), *Biologia Gallo Helvetica*, 12, 273-281.
- Ustaoglu, M. R., 1989.** Marmara Gölü' nün (Salihli) Zooplanktonu Üzerine Araştırmalar Ege Üniversitesi Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, *Doktora Tezi*, 162s.
- Ustaoglu, M. R. and Balık, S., 1990.** Zooplankton of Lake Gebekirse. *Rapp. Comm. Int. Mer. Medit*, 32(1), 23.
- Ustaoglu, R. M., 2004.** Türkiye İçsuları Zooplankton Kontrol Listesi. *Ege Üniversitesi Su Ürünleri Dergisi*, 21(3-4), 191-199.
- Yıldız, Ş., Altındağ, A., and Ergönül, M.B., 2007.** Seasonal Fluctuations in the Zooplankton Composition of a Eutrophic Lake: Lake Marmara (Manisa, Turkey), *Turkish Journal of Zoology*, 31, 121-126.
- Yiğit, S., 2002.** Seasonal Fluctuation in the Rotifer Fauna of Kesikköprü Dam Lake (Ankara, Turkey). *Turkish Journal of Zoology*, 26, 341-348.
- Yiğit, S., 2006.** Analysis of the Zooplankton Community by the Shannon-Weaver Index in Kesikköprü Dam Lake, Turkey, *Ankara Üniversitesi Ziraat Fakültesi Tarım Bilimleri Dergisi*, 12(2), 216-220.
- Yiğit, S. and Altındağ, A., 2005.** The zooplankton fauna and seasonal distribution of Beyşehir Lake, *Gazi Üniversitesi, Gazi Eğitim Fakültesi Dergisi*, 24, 217-225.