

## Estimation of growth parameters and mortality rate of *Upeneus sulphureus* (Cuvier, 1829) in the Persian Gulf ecosystem

Vahabnezhad A.<sup>1\*</sup>; Taghavimotlagh S.A.<sup>1</sup>; Salarpouri A.<sup>2</sup>

Received: October 2019

Accepted: April 2020

### Abstract

Population parameters were estimated for goatfish (*Upeneus sulphureus*, Cuvier, 1829) as a small size and abundant species in the Persian Gulf ecosystem. A method of Tropfish R package based on length frequencies analysis was used. A total of 520 fish samples were taken monthly from landing stations in the coastal area of Iranian Waters in the Persian Gulf (Bushehr coasts) from January 2017 to December 2018. The length (FL) of the captured fishes varied from 8.5 to 17 cm with an average ( $\pm$ SD) of 11.20 ( $\pm$ 2.24) cm.

The values of the von Bertalanffy growth function fit length frequency data for *U. sulphureus* (combined sex) were estimated, as  $k = 1.2$  per year,  $L_{\infty} = 19.43$  cm ( $L_F$ ). Total, natural and fishing mortality calculated, as  $Z = 3.74$  per year,  $M = 2.21$  per year and  $F = 1.53$  per year, respectively. The exploitation ratio (E) was estimated at 0.4.

The length-weight relationships of both sexes was estimated as,  $\text{Log } M_B = -1.707 + 3.00 \log L_F$  ( $r = 0.92$ ) which exhibited positive isometric growth. According to results, it could be concluded that goatfish (*Upeneus sulphureus*) is a relatively fast-growing and short-lived species. Although this species is not a direct target of fishermen, its ecological role in the Persian Gulf's food web due to their high abundance and high growth rate is undeniable.

**Keywords:** Growth, mortality rate, *Upeneus sulphureus*, Persian Gulf

1- Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran.

2- Persian Gulf and Oman Sea Ecological Research Center, Iranian Fisheries Science Research Institute, Agricultural Research, Education and Extension Organization, Bandar Abbas, Iran.

\*Corresponding author e-mail: avn9400@gmail.com.

## Introduction

*Upeneus sulphureus* (Mullidae family) is an endemic, small-sized fish which very abundant in the Persian Gulf ecosystem and commonly known as 'goatfish'. This species are distributed worldwide in tropical, subtropical and temperate waters (Lewis and Pring, 1986). Goatfishes distributed in a wide range of habitats and are associated to the bottom of the littoral, but some species of the *Upeneus* could be seen in deeper areas (Randall, 2004). All species of family Mullidae are benthic carnivores that 18 species of them have been detected in the Persian Gulf and Oman Sea and constitute one of the most important component of fish bycatch in the region (Kamrani and Daliri, 2018).

More than 90% of Iranian fishing in the Persian Gulf and Oman Sea comes from small scale fisheries (Shahifar, 2018). According to a study conducted by Hoveizavi *et al.* (2016), although the amount of fish bycatch landed by each vessel in the northwest of the Persian Gulf was low, but the high number of fishing vessels operating in this part of the Persian Gulf cause the high number of total fish bycatch, The study conducted by Eskandari *et al.* (2016) in the northwest of the Persian Gulf, showed that the highest amount of bycatch species in terms of weight per unit of effort were Belonging to *U. sulphureus* (18.41%) and the lowest value were belonging to Apogon family (0.03%). According to various studies, the ecological role of Goatfish in the food chain of the Persian Gulf is very

important. Based on the Vahabnezhad (2015), most of the stomach contents of predatory fishes, like *Argyrops spinifer*, *Pomadasys kaakan* and *Trichiurus lepturus*, *Nemiopterus japonicus* and *Saurida tumbile*, consist of Goatfish species.

Additionally, Taghavi Motlagh (2019) pointed out that the composition of the catches in the coastal area of the Persian Gulf and Oman Sea are changed from high-value species to lower value species during the past two decades which might be attributed to overexploitation of high-value species with higher mean trophic level. Also, the results of fish survey (swept area method) in the Persian Gulf and Oman Sea revealed that the CPUA for Mullidae family had increased from 585 tons in 2003 to 5926.2 tons in 2018 (90 percent increase) (Valinassab *et al.*, 2019).

Many studies have been conducted on biology and population dynamics of goatfish in different marine regions of the world, but despite a relatively high abundance of this species in the Persian Gulf and Oman Sea, no studies have been carried out on this species in Iranian part of these two ecosystems. In the northern Persian Gulf, Iraqi waters, Mohamed and Saleh (2000) have been studied some aspects of the biology of goatfish.; Federizon (1993) in Philippines waters, Mohamed *et al.* (2004); Silvestre and Garces (2004); İşmen (2005); Mutlak and Mohamed (2006); Yousif *et al.* (2007); Mohamed and Resen (2010); Gumanao *et al.*

(2016); and Kochi (2017) in the Indian ocean.

Length frequency data were used to estimate the population parameters to fill the information gap on this species. As the goatfish is an abundant fish species in the Persian Gulf, the results of the present study could be used for further ecological analysis of its role in food web structure of the Persian Gulf.

### Material and methods

The samples of *U. sulphureus* were collected from January 2017 to

December 2018 (monthly) from Persian Gulf around Bushehr coast by weir locally named Gagoor and gill nets. (Fig. 1). Totally, the fork length of 520 individuals were measured randomly during seven months to estimate the size distribution, growth parameters, and mortality rates. During sampling events, the fork length ( $L_F$ , 1 cm), body weight ( $B_W$ , 0.1g) were measured.




Figure 1: Map of the Persian Gulf showing fish landing sites along coasts (red dots).

### Growth parameter

TropFishR (Mildenberger *et al.*, 2017), an R package ver. 3.5.1 for tropical fisheries analysis, was used to estimate growth and mortality parameters.

TropFishR has enhanced functions of the FAO-ICLARM Stock Assessment Tools II FISAT II (Gayanilo *et al.*, 2005b). Monthly length-frequency data of the *U. sulphureus*, classified into 11

length classes of 0.5 intervals, were obtained from `lfq Create` function.

The bootstrapped ELEFAN with Simulated annealing (ELEFAN\_SA) function (Mildenberger *et al.*, 2017) was used for estimation of parameters  $L_{\infty}$  and  $K$  of the von Bertalanffy equation. The simulated annealing algorithm begins with a highly randomized search, which gradually focuses on regions of the parameter search space with the highest  $R_n$  scores.

Growth was modeled based on von Bertalanffy growth function (VBGF) (Sparre and Venema, 1992):

$$L_t = L_{\infty} (1 - \exp^{-K(t-t_0) + S(t) - S(t_0)})$$

Where  $L_t$  is length-at-age  $t$ ,  $L_{\infty}$  is asymptotic length,  $K$  is the von Bertalanffy growth constant, and  $t_0$  is the theoretical age when length equals zero. Usually  $t_0$  is negative, resulting in a positive length at the time of recruitment ( $t = 0$ ).  $S(t) = (CK/2\pi) \sin 2\pi(t - t_s)$ ,  $C$  is a constant indicating the amplitude of the oscillation, typically ranging from 0 to 1 (a value  $>1$  implies periods of shrinkage, which is rare), and  $t_s$  is the fraction of a year (relative to the age of recruitment,  $t=0$ ) where the sine wave oscillation begins (i.e. turns positive). In TropFishR, these additional oscillation parameters are written as  $C$  and  $t_s$ . A seasonally oscillating

VBGF was used to assess the growth parameters, which are attributed to changes in water temperature, precipitation and/or to the availability of food (Morales-Nin and Panfili, 2005).

The  $L_{\infty}$  and  $K$  were used to calculate the growth performance index ( $\Phi'$ ) =  $\log K + 2 \log L_{\infty}$  (Pauly and Munro, 1984).

The parameter  $t$  anchor indicates the fraction of the year where yearly repeating growth curves cross length equal to zero.  $T_{\max}$  derived from Pauly *et al.* (1996) formula:  $T_{\max} = 3/K + t_0$ .

#### *Mortality and exploitation rate*

The instantaneous total mortality coefficient ( $Z$ ) was estimated using Powell-Wetherall method ( $L^- = a + bL^-$ ), where  $L^-$  is the mean length and  $L'$  is the lowest length class of the collected samples, from which  $Z/K$  could be calculated from  $Z/K = -1 - b/b$  equation. This method allows estimation of  $L_{\infty}$  and  $Z/K$  from a sample representing a steady-state population, as can be approximated by pooling a time series of length-frequency data (Pauly, 1990). The rate of natural mortality ( $M$ ) was estimated according to the empirical equation as "Pauly\_Linf" function (Pauly, 1990):

$$\log_{10}(M) = -0.0066 - 0.276 \log_{10}(L_{\infty}) + 0.6543 \log_{10}(K) + 0.4634 \log_{10}(T)$$

Where  $T$  is the annual mean of sea surface temperature (*i.e.*, 26.5°C in the Persian Gulf (Ebrahimi *et al.*, 2005)). Following the calculation of  $Z$  and  $M$ , fishing mortality ( $F$ ) estimated using the functions;  $F = Z - M$  and exploration rate ( $E$ ) estimated using  $F/Z$  function (Sparre and Venema, 1992).

#### *Length-weight relationship*

The relationship between length and body mass for both sex was calculated using:  $M_B = aL_F^b$ , where  $a$  is scaling constant and  $b$  is allometric growth parameter (Hayes *et al.*, 1995).

Using the linear regression of the log-transformed equation:  $\log(M_B) = \log(a) + b \log(L_F)$ , the parameters  $a$  and  $b$  were calculated with ' $a$ ' representing the intercept and ' $b$ ' the slope of the relationship. The significant difference of  $b$  values from the isometric threshold, *i.e.*, 3, was tested by the t-test (Pauly 1984).

## **Result**

### *Growth parameters*

Mean fork length ( $\pm$  SD) of the goatfish in the Persian Gulf over the study period was recorded  $11.20 \pm 2.24$  cm. The weight of fish samples was ( $N=520$ ) varied between 11.73 to 91.89 g with the average of  $29.19 \pm 10.88$  g.

Figure 2 showed restructure of length-frequency histogram of *U. sulphureus* in the Persian Gulf. The following plot shows reconstructed frequencies, with negative and positive values as white and black colored histograms, respectively. The sum of the positive


peaks were indicated with black bars (ASP) equal to 0.16. Three age groups (cohorts) were observed, the smallest one was in December 9-12 cm ( $MA = 7$ ).

Figure 3 indicates seasonal oscillations of the growth rate. The parameter  $c$ , the "amplitude", takes value between 0 and 1. When inserting parameter  $c$  equal to 0 ( $c=0$ ), the seasonality oscillations didn't appear in the von bertalanffy growth rate, but when  $c$  equal 1 ( $c=1$ ), used the seasonal oscillations more pronounced, and growth rate appear to be zero (curve is horizontal) at the winter points ( $t_w$ ) in 0.7 and 1.7 year.


The results of the Simulated annealing (ELEFAN\_SA) analysis provided the best fit of growth model as,  $L_\infty = 19.43$  cm and  $K = 1.2$  per year (Table 1 and Fig. 4). The values of  $t_0$  and  $\phi'$  were estimated as -0.15 year, and 2.65, respectively. Ageing analysis showed that the lifespan of goatfish in this area may reach 2.35 years.

### *Mortality rate*

The natural mortality ( $M$ ) estimated by Pauly's method, as  $2.21 \text{ yr}^{-1}$ . The annual rate of total mortality ( $Z$ ) was derived from Power-Wetheral method as  $3.74 \text{ year}^{-1}$  (Fig. 5).



**Figure 2:** Restructure of length-frequency histogram of *U. sulphureus* in the Persian Gulf (2017-2018), the black and white bars are positive and negative deviations from the ‘weighted’ moving average.



**Figure 3:** The seasonlaized von Bertalanffy growth of *U. sulphureus*. Note for  $c=1$  (blue line) the growth rate is zero at the winter points.

**Table1: Summary of ELEFAN\_SA analysis for *u. sulphureus***

Asymptotic length ( $L_{\infty}$ )	Growth rate (K)	$t_{\text{anchor}}$	Phil	Rn_max
19.43	1.2	0.14	2.47	0.52




Figure 4: Refined searches using optimization algorithms with ELEFAN\_SA analysis.




Figure 5: Estimation of total mortality using Powell-Wetherall method for *U. sulphureus*.

Using the value of  $Z$  and  $M$ , fishing mortality were calculated  $1.53 \text{ year}^{-1}$ . The exploitation ratio ( $E$ ) was estimated as 0.4.

#### Length-weight relationship

The length-weight relationship and correlation coefficient ( $r$ ) for both together presented in (Fig. 6).

The value of 'a' and 'b' were -1.707 and 3.00 respectively for both sex. The b value was not significantly different from 3 (t-test;  $p > 0.05$ ). The equation of the length-weight relationships were stated below:

$$\text{Log } M_B = -1.707 + 3.00 \log L_F \quad (r = 0.92)$$




Figure 6: Length-weight relationship of *U. sulphureus* in the Persian Gulf.

#### Discussion

In the present study Length measurement has been used to estimate the key parameters of stock assessment and its population parameters like asymptotic length ( $L_{\infty}$ ), growth coefficient ( $K$ ), total mortality ( $Z$ ), natural mortality ( $M$ ) and fishing mortality ( $F$ ). Length frequency data provide unique and valuable information for modeling growth (Laslett *et al.*, 2004). In this study, length frequency data of goatfish in the Persian Gulf (Bushehr coastal water) were used to depict its growth model. The present study revealed that the highest growth oscillation occurred in January, with two peaks that each peak represents an age group. This could be due to active feeding before spawning

time (January–March) according to Akter *et al.* (2020) reports. the von bertalanffy curve which does not reveal the oscillation of the growth, The oscillation curve shows the point where the growth of the fish is stop, which in the present study, showed that according to oscillation curve when we use  $c=1$  it shows that growth of goatfish stop in age 0.7 and 1.7 years. The highest frequency was observed for length interval of 11-12 cm.

$K$  is a curvature parameter that represents how fast the fish approaches its  $L_{\infty}$ . Based on calculated  $K$  (1.2 per year), goatfish could be classified as short-lived species which reaches  $L_{\max}$  (18.50 cm) after 2 years. Table 2, present the comparison of the von Bertalanffy growth parameter estimated



in the present study with previous investigations, which shows variability of growth parameters of Goatfish in different regions which could be

attributed to various ecological condition of each ecosystem (Sparre and Venema, 1992).

**Table 2: Growth parameters of *U. sulphureus* in various areas.**

Reference	Linf	K	t <sub>0</sub>	b	a	Ø	M	Area
Present study	19.43	1.2	-0.15	3.00	0.01963	2.65	2.21	Persian Gulf
(Rajkumar, 2004)	24.5	0.67	-0.14	-	-	2.60	4.06	Visakhapatnam
(Gumanao <i>et al.</i> , 2016)	23.5	1.30	-	3.68	0.00550	2.85	-	Philippines
(Silvestre and Garces, 2004)	17	0.85	-	3.035	0.820	2.39	2.21	Philippines
(Kochi, 2017)	20.2	0.97	-	-	-	2.60	-	India
(Loychuen, 2010)	21.5	1.01	-	-	-	2.67	-	Thailand
(Chan and Liew, 1986)	23.0	1.01	-	-	-	2.76	-	Malaysia
(Federizon, 1993)	-	-	-	3	0.01930	-	-	Philippines
(Mohamed and Resen, 2010)	18.82	0.75	-	3.01	0.0621	2.36	1.85	Northwest Persian Gulf
(Prayitno <i>et al.</i> , 2020)	-	-	-	3.12	0.0159	-	-	Java Sea

The value of growth performance index ( $\dot{O}$ ) (2.65) in the present study was similar to many previously published reports in Fish Base (Table 2). According to Gayanilo *et al.* (2005b), the growth performance index ( $\dot{O}$ ) is a species-specific parameter, i.e. its values are usually similar within related taxa and have small distributions. growth dissimilarity of  $\dot{O}$  for a number of stocks of the same species or related species is an indication of the unreliability in the accuracy of estimated growth parameters. It must be realized however, that  $\dot{O}$  can be used only to compare the growth performance of fish with similar shapes (Gayanilo *et al.*, 2005a). Preliminary analyses of Moreau *et al.* (1986) suggest that the C.V. of  $\dot{O}$  for several stocks of the same species should not exceed 5 percent, which may provide some guideline that values of  $\dot{O}$  are

credible or not (Gayanilo and Pauly, 1997). The coefficient of variation of 6.41% together with other measures of dispersion (range = 0.49, variance = 0.03, standard deviation = 0.16 and mean = 2.61) for  $\dot{O}$  values presented in Table 2.

The b value of length-weight relationship for *U. sulphureus* was similar to those reported by other authors; like Mohamed and Resen (2010) and Federizon (1993).

The instantaneous rates of total (Z) natural (M) and fishing mortalities were 3.74, 2.21 and 1.53 year<sup>-1</sup>, respectively. The value of natural mortality to growth coefficient (M/K) of *U. sulphureus* in the present study was 1.84. Beverton and Holt (1957) pointed out that the value of M/K is ranging from 1.5 to 2.5 and the best value is equal to 2.0. Any deviation from this value, the fish will be vulnerable and exposed to high

natural mortality before reaching to a certain age (Gayanilo and Pauly, 1997).

The ratio of F/Z or exploitation ratio is a measure of the intensity by which a fish stock is exploited. The exploitation ratio (E) for this species in the Persian Gulf ecosystem was estimated to be 0.40, which was less than the optimum value of 0.50 suggested by Gulland (1971). The values of F and E obtained for *U. sulphureus* in the present study was similar to those reported by Mohamed and Resen (2010) in the Persian Gulf, whereas higher natural mortality rate was estimated by present study.

According to results of the Swept area surveys project, the abundance of Mullidae stock increased 90 % in past two decades in the Persian Gulf and the Oman Sea (Valinassab *et al.*, 2019) which shows that the population of small-demersal fish species are become more abundant, which could be attributed to overexploitation of top predator in this area.

As many of the small demersal species in the bottom-trawl surveys in the Persian Gulf and Oman Sea was not recorded during survey period to see the abundance variation of these species group, therefore it is suggested to record the abundance and catch per unit area of all small demersal fish species, to quantify and trace the fluctuations of the abundance of this species in this region could draw out indicators to figure out the changes occurred in this ecosystem (Ouellet *et al.*, 2016).

## References

- Akter, M., Sharifuzzaman, S., Shan, X. and Rashed-Un- Nabi, M., 2020.** Reproduction, growth, mortality and yield of the goatfish (*Upeneus sulphureus*) off Northern Bay of Bengal. *Bangladesh Journal of Ichthyology* in press.
- Beverton, R. and Holt, S., 1957.** On the dynamics of exploited fish populations. Fisheries Investigations Series II. Marine Fisheries, Great Britain Ministry of Agriculture, Fisheries and Food 19.
- Chan, E. and Liew, H., 1986.** Characteristics of an Exploited Tropical Shallow-Water Demersal Fish Community in Malaysia.
- Ebrahimi, M., Mortazawi, M., Ejlali, K., Aghajari, N., Jokar, K., Akbarzadeh, G., Seraji, F. and Aghajari, S., 2005.** Hydrology and hydrobiology studies on the Persian Gulf in Hormozgan, south of Iran (in Persian) ed. Iranian fisheries science research institute, Iran, 130P.
- Eskandari, G., Koochaknejad, E., Mayahi, Y. and Ansari, H., 2016.** Rate, ratio and amount of annual discards in commercial trawl net in northwestern part of the Persian Gulf (Khuzestan coastal waters). *Journal of Marine Science and Technology*, 15, 84-99.
- Federizon, R., 1993.** Using vital statistics and survey catch composition data for tropical multispecies fish stock assessment: application to the demersal resources

- of the central Philippines. Verlag nicht ermittelbar.
- Gayanilo, J. and Pauly, D., 1997.** FAO-ICLARM stock assessment tools (FiSAT): Reference Manual, vol. 8. FAO Computerized Information Series (Fisheries).
- Gayanilo, F.C., Sparre, P. and Pauly, D., 2005a.** FAO-ICLARM stock assessment tools II: User's guide. Food & Agriculture Org.
- Gayanilo, F.C., Sparre, P. and Pauly, D., 2005b.** FAO-ICLARM stock assessment tools II: User's guide. Food & Agriculture Org.
- Gulland, J.A., 1971.** Fish resources of the ocean.
- Gumanao, G., Saceda-Cardoza, M., Mueller, B. and Bos, A., 2016.** Length–weight and length–length relationships of 139 Indo-Pacific fish species (Teleostei) from the Davao Gulf, Philippines. *Journal of Applied Ichthyology*, 32, 377-385.
- Hoveizavi, S., Doustshenas, B., Savari, A., Eskandari, G., Mohammadasgari, H. and Jamali, H., 2016.** Length-Weight Relationships for Eight Species of By-Catch and Discard Fishes in the Fishing Grounds of Khuzestan Coastal Waters (Northwest Persian Gulf). *Advances in BioResearch* 7.
- İşmen, A., 2005.** Age, growth and reproduction of the goldband goatfish, *Upeneus moluccensis* (Bleeker, 1855), in Iskenderun Bay, the Eastern Mediterranean. *Turkish Journal of Zoology*, 29, 301-309.
- Kamrani, E. and Daliri, M., 2018.** ical Characteristics of *Upeneus doriae* (Günther, 1869) in the coast of Bandar Abbas. *Journal of Applied Ichthyological Research*, 6, 13-30.
- Kochi, C., 2017.** CMFRI annual report 2016–2017. Technical Report CMFRI, Kochi.
- Laslett, G.M., Eveson, J.P. and Polacheck, T., 2004.** Fitting growth models to length frequency data. *ICES Journal of Marine Science*, 61, 218-230.
- Lewis, A.D. and Pring, C.K., 1986.** Freshwater and brackishwater fish and fisheries of Fiji In T. Petr (ed.) Reports and papers presented at the Indo-Pacific Fishery Commission Expert Consultation on inland fisheries of the larger Indo-Pacific islands. Bangkok,. 258
- Loychuen, K., S. Hoimuk, S. Boonsook, J. and Ritthisaman, S., 2010.** tock assessment of sunrise goatfish *Upeneus sulphureus* (Cuvier, 1829) along the Andaman Sea Coast of Thailand, 41.
- Mildenberger, T.K., Taylor, M.H. and Wolff, M., 2017.** TropFishR: an R package for fisheries analysis with length-frequency data. *Methods in Ecology and Evolution*, 8, 1520-1527.
- Mohamed, A. and Saleh, J., 2000.** Some aspects of the biology of *Upeneus sulphureus* in the northwest Arabian Gulf, Iraq. *Marina Mesopotamica*, 15, 581-592.
- Mohamed, A., Hussein, S. and Mutlak, F., 2004.** Food habit of *Upeneus sulphureus* in Iraqi marine waters, north west Arabian

- Gulf/Iraq. *Marina Mesopotamica*, 19, 171-187.
- Mohamed, A. and Resen, A., 2010.** The population status of sulphur goatfish *Upeneus sulphureus* in the Iraqi marine waters, northwest Arabian Gulf. *Mesopotamian Journal of Marine Science*, 25, 31-40.
- Morales-Nin, B. and Panfili, J., 2005.** Seasonality in the deep sea and tropics revisited: what can otoliths tell us? *Marine and Freshwater Research*, 56, 585-598.
- Moreau, J., Bambino, C., Pauly, D., Maclean, J., Dizon, L. and Hosillos, L., 1986.** The first Asian fisheries forum, The first Asian fisheries forum. Asian Fish. Soc.
- Mutlak, F. and Mohamed, A., 2006.** Monthly fluctuations of catch rates of three goatfishes species from Khor Al-Amayh, Northwest Arabian Gulf. *Mesopotamian Journal of Marine Science*, 21, 257-270.
- Ouellet, P., Savenkoff, C., Benoît, H.P. and Galbraith, P.S., 2016.** A comparison of recent trends in demersal fish biomass and their potential drivers for three ecoregions of the Gulf of St Lawrence, Canada. *ICES Journal of Marine Science*, 73, 329-344.
- Pauly, D., 1990.** Length-converted catch curves and the seasonal growth of fishes. *Fishbyte*, 8, 33-38.
- Pauly, D., Moreau, J. and Gayanilo Jr, F., 1996.** A new method for comparing the growth performance of fishes, applied to wild and farmed Tilapias, The third international symposium on tilapia in aquaculture. ICLARM Conf. Proc. 41, 433-441.
- Pauly, D. and Munro, J., 1984.** Once more on the comparison of growth in fish and invertebrates. *Fishbyte*, 2, 1-21.
- Prayitno, M., Setiawan, H., Jatmiko, I., Rahman, M. and Wiadnya, D., 2020.** Spawning potential ratio (SPR) of Sulphur Goatfish (*Upeneus sulphureus*): biological basis for demersal fishery management in Java Sea.
- Rajkumar, U., 2004.** Fishery and some aspects of population dynamics of goatfish *Upeneus vittatus* (Forsk.) off Visakhapatnam. *Journal of the Marine Biological Association of India*, 46, 192-197.
- Randall, J.E., 2004.** Revision of the goatfish genus *Parupeneus* (Perciformes: Mullidae) with descriptions of two new species. Bishop Museum.
- Shahifar, R., 2018.** Iran tuna fisheries Sharks by-catch in IOTC area of competence in 2017. WPEB14, South Africa, 10-14 Sept 2018. IOTC-2018-WPEB14-17\_Rev1. 5P.
- Silvestre, G.T. and Garces, L.R., 2004.** Population parameters and exploitation rate of demersal fishes in Brunei Darussalam (1989–1990). *Fisheries Research*, 69, 73-90.
- Sparre, P. and Venema, S., 1992.** Introduction to tropical fish stock assessment part 1. manual FAO FISH. Tech. Pap 306.

- Taghavi Motlagh, S., 2019.** The analysis of catches data and the forecast of the production of species in the southern waters of the country, based on catch data series (2014-1990). Iranian Fisheries Science Research Institute, Iran, 450.
- Vahabnezhad, A., 2015.** Feeding habits and trophic levels of some demersal fish species in the Persian Gulf (Bushehr Province) using Ecopath model. Islamic Azad University, Science and Research Branch, Tehran.
- Valinassab, T., Dehghani, R., Kamali, A., Behzadi, S., Darvishi, M. and Salarpour, A., 2019.** Biomass estimation of demersal fishes by swept area method in the Persian Gulf and Oman Sea. Iranian Fisheries Science Research Institute, Iran, 349.
- Yousif, U., Hussain, N. and Mohamed, A., 2007.** The effectiveness of the carnivorous fishes on food resources in the Northwest Arabian Gulf. *Mesopotamian Journal of Marine Science*, 22, 25-36.