

Iman Adomian decomposition method applied to Logistic differential model

I. A. Almardy

Department of Management Information Systems and Production Management, College of Business and Economics, Qassim University, P.O.Box: 6640, Buraidah 51452, Saudi Arabia

Nagat A. A. siddig

Department of Mathematics, College of Sciences and Arts, King Kalied University, Dhahran Aljonoub.

Samah A. H. Fodol

Department of Mathematics, College of Sciences, King Kalied University.

H. A. Albushra

Department of Electrical Engineering College of Engineering and Information Technology, Buraidah private Colleges.

A. K. Osman

Department of Management Information Systems and Production Management, College of Business and Economics, Qassim University, P.O.Box: 6640, Buraidah 51452, Saudi Arabia

Abstract

This study applies Iman Adomian Decomposition Method (IADM) to solve the Logistic Differential Model (LDM) of different forms and coefficients. Illustrative examples are considered, and the obtained results are in good agreement compared to those already in the literature. This study, therefore, recommends the proposed method (IADM) for application in other aspects of applied mathematics for real-life problems.

Keywords: *Logistic differential model, non-linear model, approximate solutions; Transform method.*

1. Introduction

A logistic differential equation is a conventional mathematical expression whose solution is a logistic function. Exponential functions fail to consider limitations that prevent infinite resources, whereas logistic functions do [1-3]. Many other areas, such as machine learning, chess ratings, cancer therapies (such as the modeling of tumor development), economics, and language

adoption studies, rely on these types of models [4, 5]. This model is unrealistic since the environment constrains population expansion.

$$\frac{dp}{dt} = rp \left(1 - \frac{p}{k}\right), p = p(x, t) \quad (1.1)$$

$$p_0 = p(x, 0)$$

where $p=p(x,t)$ is the population size of the species at time t , r , denotes the rate of growth in the absence of limited resources, and k denotes the carrying capacity or the maximum

population that the ecosystem can support indefinitely.

The goal of this study is to apply the AADM to find a solution for the Logistic Differential Model (LDM) [6-10]. The aim is to present a simple and practical method for obtaining a better approximation to find the exact solution to the LDM. Thus, the objectives are to: apply AADM to the logistic differential equation and compare the results obtained via applying the AADM and exact solutions of the Logistic differential model. Although, numerical approaches have been arbitrated more efficient and reliable in solving the dynamical models (equations) and other differential models in this regard [11-20].

Different solution experts have recently discussed numerous methods for finding an exact or numerical solution to ordinary or partial differential models [21-26]. In this work, a novel approach termed Successive Approximation Method (SAM) is applied to some non-linear evolution models.

2. Note on ImanAdomian Decomposition Method (IADM)

The AADM will be discussed here in relation to the Logistic Differential Equation.

2.1 Adomian Decomposition Method (ADM)

Let us examine the differential equation of the following form:

$$Dw + Rw + Nw = g(x, t), w = w(x, t) \quad (2,1)$$

where the linear operator (differential) is D , the differential operator has a remaining part R and a nonlinear, while $g = g(x, t)$ is a source term. Generally, we choose $D = \frac{d^n}{dx^n}(\cdot)$, to be the n th-order differential operator, has its inverse D^{-1} follows as the n th-order integration

operator. Therefore, the inverse linear D^{-1} used on (2.1), we have

$$D^{-1}[Dw + Rw + Nw] = D^{-1} g(x, t) \quad (2,2)$$

Where,

$$D^{-1} Dw = y - \alpha \quad (2,3)$$

And α signifies the initial value. α

Thus, (2.3) becomes:

$$y - \alpha + D^{-1}[Rw + Nw] = D^{-1}g \quad (2,4)$$

$$y = D^{-1}g + \alpha - D^{-1}[Rw + Nw] \quad (2,5)$$

$$y = \beta(y) - D^{-1}[Rw + Nw] \quad (2,6)$$

Where

$$\beta(y) = D^{-1}g + \alpha \quad (2,7)$$

which signifies a function obtained by integrating the source term with respect to the initial condition(s). The ADM expresses the solution $y(t)$ in the series form:

$$y = \sum_{n=0}^{\infty} y_n \quad (2,8)$$

Also, the non-linear component can be stated as:

$$Nw(x, t) = \sum_{n=0}^{\infty} A_n \quad (2,9)$$

$$A_n = \frac{1}{n!} \frac{d^n}{d\lambda^n} \left(f \left(t, \sum_{k=0}^{\infty} \lambda^k y_k \right) \right)_{\lambda=0}, n \geq 0 \quad (2,10)$$

$$\sum_{n=0}^{\infty} y_n = \beta(y) - D^{-1} \left[R \left(\sum_{n=0}^{\infty} y_n \right) + \sum_{n=0}^{\infty} A_n \right] \quad (2,11)$$

By a recursive equation, we have:

$$y_0(x) = \beta(x) \quad (2,12)$$

$$y_{n+1}(x) = -D^{-1}[Ry_n + A_n], n \geq 0 \quad (2,13)$$

Thus, the solution is:

$$y(x) = \lim_{n \rightarrow \infty} \left(\sum_{n=0}^{\infty} y_n \right) \quad (2,14)$$

2.2 Iman Transform Method (ITM)

The Imantransform method helps in solving differential equations (ODE) and partial (PDE) in the time domain. It is also used as an effective tool in response to initial data analyze the fundamental properties of a linear system governed by the differential equation.

2.3. Definition of Iman Transform

Let C a function such that

$$C = \{H(t): |H(t)| < Me^{t|k|}, \text{ for } M, k_1, k_2 > 0\} \quad (2,15)$$

Thus, the Iman transform of H(t) is defined and denoted as:

$$I[H(t)] = H(v) = \frac{1}{v^2} \int_0^{\infty} H(t) e^{-tv^2} dt \quad (2,16)$$

2.4 Properties of Iman Transform

The main properties of Iman Transform are:

$$PI1: I[1] = \frac{1}{v^4}$$

$$PI2: I[t] = \frac{1}{v^6}$$

$$PI3: I[e^{at}] = \frac{1}{v^4 - av^2}$$

$$PI4: I[t^n] = \frac{n!}{v^{2n+4}}$$

2.5 ImanAdomian Decomposition Method (IADM)

The IADM consists of a mix of both the Iman transform method and the Adomian decomposition approach. The problem can either be linear or non-linear.

Let us consider the general differential equation of the form:

$$Dw + Rw + Nw = g(x, t), w = w(x, t) \quad (2,17)$$

where D, N, R and g are as defined earlier.

Suppose

$$g(x, 0) = g_1,$$

$$w = w(x, t)$$

then the Iman transform of (2.17) is as follows:

$$I(Dw) + I(Rw) + I(Nw) = I(g)$$

$$I(Dw) = I(g) - I(Rw) - I(Nw)$$

$$v^2 I(x, v) - \frac{w}{v^2} = I(g) - I(Rw + Nw)$$

$$I(x, v) = \frac{w}{v^4} + v^2 I(g) - v^2 I(Rw + Nw)$$

$$I(x, v) = G(x, t) - v^2 A(Rw + Nw) \quad (2,18)$$

Where $G(x, t)$ is the resulting term from the source and initial condition terms when used.

Based on the inverse Iman transform of (2.18), we have:

$$I^{-1}I(x, v) = I^{-1}[G(x, t)] - I^{-1}[v^2 I(Rw + Nw)]$$

$$h = I^{-1}[G(x, t)] - I^{-1}[v^2 A(Rw + Nw)] \quad (2,19)$$

Using ADM, the series solution is defined as

$$w = \sum_{n=0}^{\infty} w_n \quad (2,20)$$

And the non-linear term as:

$$\begin{cases} Nw = \sum_{n=0}^{\infty} A_n \\ A_n, \text{ as Adomian polynomials.} \end{cases} \quad (2,21)$$

Hence, (2.21) becomes

$$\begin{cases} w_1 = g_1 * \\ w_{n+1} = I^{-1}[v^2 I(R(w_n) + (A_n))] \end{cases} \quad (2,22)$$

3. Method and Model Discussed

This part discusses the proposed method and the Logistic model, as formulated based on some assumptions. Case examples are also considered via the IADM.

CASE I: Consider the following version of the LDE:

$$\left\{ \frac{dp}{dt} = \frac{1}{4}p(1-p), \quad p(0) = \frac{1}{3} \right. \quad (3,1)$$

whose exact solution is:

$$p(t) = \frac{e^{0.25t}}{2 + e^{0.25t}} \quad (3,2)$$

By the IADM, we have

$$I[P_t] = I\left[\frac{P}{4} - \frac{P^2}{4}\right] \quad (3,3)$$

$$v^2 l(x, v) - \frac{P(x, 0)}{v^2} = I\left[\frac{P}{4} - \frac{P^2}{4}\right] \quad (3,4)$$

$$l(x, v) = v^2 I\left[\frac{P}{4} - \frac{P^2}{4}\right] + \frac{P(x, 0)}{v^4} \quad (3,5)$$

$$l(x, v) = v^2 I\left[\frac{P}{4} - \frac{P^2}{4}\right] + \frac{1}{3v^4} \quad (3,6)$$

Taking the Iman inverse of (3.6) gives:

$$I^{-1}[l(x, v)] = I^{-1}\left[v^2 A\left[\frac{P}{4} - \frac{P^2}{4}\right]\right] + I^{-1}\left[\frac{1}{3v^4}\right]$$

$$p(x, t) = I^{-1}\left[v^2 I\left[\frac{P}{4} - \frac{P^2}{4}\right]\right] + \frac{1}{3} I^{-1}\left[\frac{1}{v^4}\right]$$

$$p(x, t) = I^{-1}\left[v^2 A\left[\frac{P}{4} - \frac{P^2}{4}\right]\right] + \frac{1}{3} \quad (3,7)$$

Next, we apply the Adomian approach to (3.7), where

$$p(x, t) = \sum_{n=0}^{\infty} p_n$$

Hence,

$$\sum_{n=0}^{\infty} p_n = \frac{1}{3} + \frac{1}{4} I^{-1}\left[v^2 I\left[\sum_{n=0}^{\infty} p_n - \sum_{n=0}^{\infty} A_n\right]\right] \quad (3,8)$$

The recursive relation is:

$$\left. \begin{aligned} P_0 &= \frac{1}{3} \\ P_{n+1} &= I^{-1}\left[v^2 I\left[\frac{p_n}{4} - \frac{A_n}{4}\right]\right], n \geq 0 \end{aligned} \right\} \quad (3,9)$$

Thus, for $n = 0, 1, 2, 3, 4, 5, \dots$, the following are respectively obtained:

$$p(t) = p_0 + p_1 + p_2 + \dots$$

$$p(t) = \frac{1}{3} + \frac{t}{18} + \frac{t^2}{432} - \frac{t^3}{5184} + \dots \quad (3,10)$$

Exact solution:

$$p(t) = \frac{e^{0.25t}}{2 + e^{0.25t}}$$

CASE II: Consider the following version of the LDE

$$\left\{ \frac{dp}{dt} = \frac{1}{2}p(1-p), \quad p(0) = \frac{1}{2} \right. \quad (3,11)$$

whose exact solution is:

$$p(t) = \frac{e^{0.25t}}{2 + e^{0.25t}} \quad (3,12)$$

By the IADM, we have

$$I[P_t] = I\left[\frac{P}{2} - \frac{P^2}{2}\right] \quad (3,13)$$

$$v^2 l(x, v) - \frac{P(x, 0)}{v^2} = I\left[\frac{P}{2} - \frac{P^2}{2}\right] \quad (3,14)$$

$$l(x, v) = v^2 I\left[\frac{P}{2} - \frac{P^2}{2}\right] + \frac{P(x, 0)}{v^4} \quad (3,15)$$

$$l(x, v) = v^2 I\left[\frac{P}{2} - \frac{P^2}{2}\right] + \frac{1}{2v^4} \quad (3,16)$$

Taking the Iman inverse of (3.6) gives:

$$I^{-1}[l(x, v)] = I^{-1}\left[v^2 I\left[\frac{P}{2} - \frac{P^2}{2}\right]\right] + I^{-1}\left[\frac{1}{2v^4}\right]$$

$$p(x, t) = I^{-1}\left[v^2 I\left[\frac{P}{2} - \frac{P^2}{2}\right]\right] + \frac{1}{2} I^{-1}\left[\frac{1}{v^4}\right]$$

$$p(x, t) = I^{-1}\left[v^2 I\left[\frac{P}{2} - \frac{P^2}{2}\right]\right] + \frac{1}{2} \quad (3,17)$$

Next, we apply the Adomian approach to (3.7), where

$$p(x, t) = \sum_{n=0}^{\infty} p_n$$

Hence,

$$\sum_{n=0}^{\infty} p_n = \frac{1}{2} + \frac{1}{2} I^{-1} [v^2 I [\sum_{n=0}^{\infty} p_n - \sum_{n=0}^{\infty} A_n]] \quad (3,18)$$

The recursive relation is:

$$\left. \begin{aligned} P_0 &= \frac{1}{2} \\ P_{n+1} &= I^{-1} \left[v^2 I \left[\frac{p_n}{2} - \frac{A_n}{2} \right] \right], n \geq 0 \end{aligned} \right\} \quad (3,19)$$

Thus, for $n = 0, 1, 2, 3, 4, 5, \dots$, the following are respectively obtained:

$$p(t) = p_0 + p_1 + p_2 + \dots$$

$$p(t) = \frac{1}{2} + \frac{t}{8} + -\frac{t^3}{384} + \dots \quad (3,10)$$

Exact solution:

$$p(t) = \frac{e^{0.25 t}}{2 + e^{0.25 t}}$$

4. Conclusions

In this work, the ImanAdomian Decomposition Method (IADM) was applied to the non-linear differential equation known as the Logistic Differential Model. The IADM has an advantage in its applicability, speed of convergence, and accuracy, unlike other numerical methods. Applying the IADM yields a series solution. The IADM is a very effective tool in the solution of the Logistic Differential Model. It can also be applied to several other more complex ordinary differential equations (both linear and non-linear). The results have shown distinctive characteristics of the method in terms of effectiveness and speed of accuracy. The IADM does not require linearization and initial guess points.

Acknowledgment

The authors extend their appreciation to the Deputyship for Research & Innovation,

Ministry of Education and Qassim University, Saudi Arabia for funding this research work through the project number (QU- 6640-51452).

Reference

- [1] Saad K. M., AL-Shomrani A. A., Mohamed S. Mohamed, Yang X. J. (2016). Solving Fractional Order Logistic Equation by Approximate Analytical Methods. Int. J. Open Problems Compt. Math., Vol. 9, No. 2, June 2016 ISSN 1998-6262.
- [2] Petropoulou N. (2010). A Discrete Equivalent of the Logistic Equation. Hindawi Publishing Corporation Advances in Difference Equations Volume 2010, Article ID 457073, 15 pages doi:10.1155/2010/457073
- [3] Vasily E. Tarasov (2020). Exact Solutions of Bernoulli and Logistic Fractional Differential Equations with Power Law Coefficients.
- [4] Lslam S., Yasir Khan, Naeem Faraz and Francis Austin (2010), Numerical Solution of Logistic Differential Equations by using the Laplace Decomposition Method, World Applied Sciences Journal 8 (9):1100-1105.
- [5] Saleh Alshammari, Mohammed Al-Smadi, Mohammad Al Shammam, Ishak Hashim and MohdAlmie Alias (2019). Advanced Analytical Treatment of Fractional Logistic Equations Based on Residual Error Functions. International Journal of Differential Equations Volume 2019, Article ID 7609879, 11 pages
- [6] Edeki S. O., Ogundile O. P., Egara F. O., Braimah J. A., Elzaki decomposition method for approximate solution of a one-dimensional heat model with axial symmetry. 2020 International Conference on Mathematics and Computers in Science

- and Engineering doi:
10.1109/MACISE49704.2020.00061
- [7] K.S.Aboodh, R.A.Farah, I.A.Almardy and F.A.Almostafa, some Application of Aboodh Transform to First Order Constant Coefficients Complex equations, International Journal of Mathematics and its Applications, ISSN : 2347-1557, App.6(1-A)(2018),1-6.
- [8] K.S.Aboodh, M.Y.Ahmed, R.A.Farah, I.A.Almardy and M.Belkhamisa, New Transform Iterative Method for Solving some Klein-Gordon Equations, (IJARSCT) IIUI, ISSN 1 Volume 2, (2022), pp.118-126. SCOPe Database Article Link: <https://sdbindex.com/documents/00000310/00001-85016.pdf>
- [8] K.S.Aboodh, I.A.Almardy, R.A.Farah, M.Y.Ahmed and R.I.Nuruddeen, On the Application of Aboodh Transform to System of Partial Differential Equations, BEST, IJHAMS Journal, ISSN(P): 2348-0521; ISSN(E): 2454-4728 Volume 10, Issue 2, Dec 2022. [[{GUIFUYFYHVHVNOOJIOHIUHG
- [9] Biazar J., Shafiof S. M. (2007). A Simple Algorithm for Calculating Adomian Polynomials. Int. J. Contemp. Math. Sciences, Vol. 2, 2007, no. 20, 975 – 982.
- [10] Nuruddeen R.I. and Nass A.M, Aboodh decomposition method and its application in solving linear and non-linear heat equations, European Journal of Advances in Engineering and Technology, 3(7) (2016) 34-37.
- [11] K.S.Aboodh, R.A.Farah, I.A.Almardy and F.A.Almostafa, Solution of partial Integro-Differential Equations by using Aboodh and Double Aboodh Transform Methods, Global Journal of pure and Applied Mathematics, ISSN 0973-1768 Volume 13, Number 8 (2017), pp.4347-4360.
- [12] Elzaki T.M. and Ezaki S.M., On the Elzaki transform and ordinary differential equation with variable coefficients, Adv. in Theo. & Appl. Math., 6 (2011) 41-46
- [13] Okoli Deborah Chikwado (2019), Solution Methods for One-Factor Bond Pricing Model, B S.c project Unpublished, Department Of Mathematics Covenant University, Ota.
- [14] Rahmatullah Ibrahim Nuruddeen (2016). Elzaki Decomposition Method and its Applications in Solving Linear and Nonlinear Schrodinger Equations. Article in Sohag Journal of Mathematics · May 2017 DOI: 10.18576/sjm/040201.
- [15] Ruchi Nigam (2015). A New Formulation of Adomian Polynomials. International Journal of Mathematics and Scientific Computing (ISSN: 2231-5330), VOL. 5, NO. 2, 2015. ISSN 2224-5804 (Paper) ISSN 2225-0522 (Online) Vol.2, No.4, 2012.
- [16] Wazwaz A.M., A reliable technique for solving linear and non-linear Schrodinger equations by Adomian decomposition method, Bulletin of institute of mathematics, 29 (2) (2001) 125-134.
- [17] Wazwaz A.M., A study on linear and non-linear Schrodinger equations by the variational iteration method, Chaos, Solitons and Fractals 37 (2008) 1136-1142.
- [18] Edeki S.O., Jena, R.M., Ogundile, O.P., Chakraverty, S. PDTM for the solution of a time-fractional barrier option Black-Scholes model, 2021, Journal of Physics: Conference Series 1734(1), 012055.
- [19] Singh J., Kumar D. and Rathore S., Application of Homotopy Perturbation Transform Method for Solving Linear and

- Nonlinear Klein-Gordon Equations, Journal of Information and Computing Science, 7 (2), (2012), 131-139.
- [20] Oghonyon J. G. , Omoregbe N. A. Bishop S.A., Implementing an order six implicit block multistep method for third order ODEs using variable step size approach, Global Journal of Pure and Applied Mathematics12 (2), 2016, 1635-1646.
- [21] Edeki S.O., P.O. Ogunniyi, O. F. Imaga, Coupled method for the solution of a one-dimensional heat equation with axial symmetry, 2021 Journal of Physics: Conference Series 1734(1),012046.
- [22] Saadatmandi A., Dehghan M., Numerical solution of hyperbolic telegraph equation using the Chebyshev Tau method, Numer. Methods Partial Differential Eq. (2009).
- [23] Mohanty R.K, Jain M.K, George K.. On the use of high order difference methods for the system of one space second order non-linear hyperbolic equations with variable coefficients, J. Comp. Appl. Math. 72 (1996), 421-431.
- [24] Oghonyon J. G. , Okunuga S. A., Bishop S. A., A 5-step block predictor and 4-step corrector methods for solving general second order ordinary differential equations, Global Journal of Pure and Applied Mathematics11 (5), 2015, 3847-386.
- [25] I. A. Almardy, S. Y. Eltayeb , S. H. Mohamed , M. A. Alkerr , A. K. Osman, H. A. Albushra, A Comparative Study of Iman and Laplace Transforms to Solve Ordinary Differential Equations of First and Second Order, ISSN 1, Volume 3, (2023), pp323-328. International Journal of Advanced Research in Science, Communication and Technology (IJARSCT).
- [26] On the Iman Transform and Systems of Ordinary Differential Equations I. A. Almardy ,R. A. Farah, M. A. Elkeer, Volume 3, Issue 1, February 2023, International Journal of Advanced Research in Science, Communication and Technology (IJARSCT), ISSN (Online) 2581-9429, Copyright to IJARSCT DOI: 10.48175/568 580 www.ijarsct.co.in.